

|  CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 123456 DATED 01/01/17  | | | | | | | | | | | | | | | | | | |
|--|--------------------------------|--------------------------------|-------------------------|----|-------|------|----|-------|------|----|--------|------|-----|------|------|-----|--------|------|
| JIM SMITH GROWER GOAT FEED MEDICATED TYPE C COMPLETE FEED CONTAINING DECCOX™(DECOQUINATE) FORMULA # JSMITHGOATJAN17 | | | | | | | | | | | | | | | | | | |
| STATEMENT OF PURPOSE | | | | | | | | | | | | | | | | | | |
| For the prevention of coccidiosis in young goats caused by Eimeria christenseni and E. ninakohlyakimovae. | | | | | | | | | | | | | | | | | | |
| ACTIVE DRUG INGREDIENT | | | | | | | | | | | | | | | | | | |
| Decoquinatone 22.73 grams/ton 11.36 milligrams/pound | | | | | | | | | | | | | | | | | | |
| USE DIRECTIONS | | | | | | | | | | | | | | | | | | |
| Feed at a rate of 22.7 mg decoquinatone per 100 lb of body weight per day as indicated in the feeding chart below. Feed at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. | | | | | | | | | | | | | | | | | | |
| <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Goat WT. (pounds)</th> <th>decoquinatone required (mg)</th> <th>lbs. To feed per day</th> </tr> </thead> <tbody> <tr><td>25</td><td>5.675</td><td>0.50</td></tr> <tr><td>50</td><td>11.35</td><td>1.00</td></tr> <tr><td>75</td><td>17.025</td><td>1.50</td></tr> <tr><td>100</td><td>22.7</td><td>2.00</td></tr> <tr><td>125</td><td>28.375</td><td>2.50</td></tr> </tbody> </table> | Goat WT. (pounds) | decoquinatone required (mg) | lbs. To feed per day | 25 | 5.675 | 0.50 | 50 | 11.35 | 1.00 | 75 | 17.025 | 1.50 | 100 | 22.7 | 2.00 | 125 | 28.375 | 2.50 |
| Goat WT. (pounds) | decoquinatone required (mg) | lbs. To feed per day | | | | | | | | | | | | | | | | |
| 25 | 5.675 | 0.50 | | | | | | | | | | | | | | | | |
| 50 | 11.35 | 1.00 | | | | | | | | | | | | | | | | |
| 75 | 17.025 | 1.50 | | | | | | | | | | | | | | | | |
| 100 | 22.7 | 2.00 | | | | | | | | | | | | | | | | |
| 125 | 28.375 | 2.50 | | | | | | | | | | | | | | | | |
| PRECAUTIONARY STATEMENTS | | | | | | | | | | | | | | | | | | |
| WARNING: Do not feed to goats producing milk for human consumption. | | | | | | | | | | | | | | | | | | |
| CAUTION: Do not use in feeds containing bentonite. | | | | | | | | | | | | | | | | | | |
| MANUFACTURED BY: BLUE BIRD FED MILL, ANY CITY, M ANY STATE 55555 Net Weight on Bag and/or Invoice | | | | | | | | | | | | | | | | | | |
| DECCOX™ is a trademark of Alpharma Inc. label revision 12/01/16 | | | | | | | | | | | | | | | | | | |

|  CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 123456 DATED 01/01/17  | | | | | | | | | | | | | | | | | | |
|--|--------------------------------|--------------------------------|-------------------------|----|-------|------|----|-------|------|----|--------|------|-----|------|------|-----|--------|------|
| JIM SMITH GROWER GOAT FEED MEDICATED TYPE C COMPLETE FEED CONTAINING DECCOX™(DECOQUINATE) FORMULA # JSMITHGOATJAN17 | | | | | | | | | | | | | | | | | | |
| STATEMENT OF PURPOSE | | | | | | | | | | | | | | | | | | |
| For the prevention of coccidiosis in young goats caused by Eimeria christenseni and E. ninakohlyakimovae. | | | | | | | | | | | | | | | | | | |
| ACTIVE DRUG INGREDIENT | | | | | | | | | | | | | | | | | | |
| Decoquinatone 22.73 grams/ton 11.36 milligrams/pound | | | | | | | | | | | | | | | | | | |
| USE DIRECTIONS | | | | | | | | | | | | | | | | | | |
| Feed at a rate of 22.7 mg decoquinatone per 100 lb of body weight per day as indicated in the feeding chart below. Feed at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. | | | | | | | | | | | | | | | | | | |
| <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Goat WT. (pounds)</th> <th>decoquinatone required (mg)</th> <th>lbs. To feed per day</th> </tr> </thead> <tbody> <tr><td>25</td><td>5.675</td><td>0.50</td></tr> <tr><td>50</td><td>11.35</td><td>1.00</td></tr> <tr><td>75</td><td>17.025</td><td>1.50</td></tr> <tr><td>100</td><td>22.7</td><td>2.00</td></tr> <tr><td>125</td><td>28.375</td><td>2.50</td></tr> </tbody> </table> | Goat WT. (pounds) | decoquinatone required (mg) | lbs. To feed per day | 25 | 5.675 | 0.50 | 50 | 11.35 | 1.00 | 75 | 17.025 | 1.50 | 100 | 22.7 | 2.00 | 125 | 28.375 | 2.50 |
| Goat WT. (pounds) | decoquinatone required (mg) | lbs. To feed per day | | | | | | | | | | | | | | | | |
| 25 | 5.675 | 0.50 | | | | | | | | | | | | | | | | |
| 50 | 11.35 | 1.00 | | | | | | | | | | | | | | | | |
| 75 | 17.025 | 1.50 | | | | | | | | | | | | | | | | |
| 100 | 22.7 | 2.00 | | | | | | | | | | | | | | | | |
| 125 | 28.375 | 2.50 | | | | | | | | | | | | | | | | |
| PRECAUTIONARY STATEMENTS | | | | | | | | | | | | | | | | | | |
| WARNING: Do not feed to goats producing milk for human consumption. | | | | | | | | | | | | | | | | | | |
| CAUTION: Do not use in feeds containing bentonite. | | | | | | | | | | | | | | | | | | |
| MANUFACTURED BY: BLUE BIRD FED MILL, ANY CITY, M ANY STATE 55555 Net Weight on Bag and/or Invoice | | | | | | | | | | | | | | | | | | |
| DECCOX™ is a trademark of Alpharma Inc. label revision 12/01/16 | | | | | | | | | | | | | | | | | | |