

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 50001 DATED 09/27/19



**Steer Dewormer CATTLE FEED
MEDICATED
FEED CONTAINING FENBENDAZOLE**

FORMULA #: Jstar Deworm 9-2016

STATEMENT OF PURPOSE

For the removal and control of lungworms (*Dictyocaulus viviparus*); barberpole worms (*Haemonchus contortus*); brown stomach worms (*Ostertagia ostertagi*); small stomach worms (*Trichostrongylus axei*); hookworms (*Bunostomum phlebotomum*); thread-necked intestinal worms (*Nematodirus helvetianus*); small intestinal worms (*Cooperia punctata and Cooperia oncophora*); bankrupt worms (*Trichostrongylus colubriformis*); and nodular worms (*Oesophagostomum radiatum*).

ACTIVE DRUG INGREDIENT

Fenbendazole 217.68 grams/ton
108.84 milligrams/pound

USE DIRECTIONS

Feed as the sole ration or as a top dress for **ONE DAY ONLY** as indicated in the chart below. Retreatment may be needed after 4 to 6 weeks.

Cattle Wt. (pounds)	Fenbendazole required (mg)	Amount to feed/day(lbs.)
200	454	4.17
400	908	8.34
600	1362	12.51
800	1816	16.69

Cattle Wt. (pounds)	Fenbendazole required (mg)	Amount to feed/day(lbs.)
1000	2270	20.86
1200	2724	25.03
1400	3178	29.20

PRECAUTIONARY STATEMENTS

WARNING: Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle, the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Net Weight on Bag and/or Invoice

revision 12/20/16

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 50001 DATED 09/27/19



**Steer Dewormer CATTLE FEED
MEDICATED
FEED CONTAINING FENBENDAZOLE**

FORMULA #: Jstar Deworm 9-2016

STATEMENT OF PURPOSE

For the removal and control of lungworms (*Dictyocaulus viviparus*); barberpole worms (*Haemonchus contortus*); brown stomach worms (*Ostertagia ostertagi*); small stomach worms (*Trichostrongylus axei*); hookworms (*Bunostomum phlebotomum*); thread-necked intestinal worms (*Nematodirus helvetianus*); small intestinal worms (*Cooperia punctata and Cooperia oncophora*); bankrupt worms (*Trichostrongylus colubriformis*); and nodular worms (*Oesophagostomum radiatum*).

ACTIVE DRUG INGREDIENT

Fenbendazole 217.68 grams/ton
108.84 milligrams/pound

USE DIRECTIONS

Feed as the sole ration or as a top dress for **ONE DAY ONLY** as indicated in the chart below. Retreatment may be needed after 4 to 6 weeks.

Cattle Wt. (pounds)	Fenbendazole required (mg)	Amount to feed/day(lbs.)
200	454	4.17
400	908	8.34
600	1362	12.51
800	1816	16.69

Cattle Wt. (pounds)	Fenbendazole required (mg)	Amount to feed/day(lbs.)
1000	2270	20.86
1200	2724	25.03
1400	3178	29.20

PRECAUTIONARY STATEMENTS

WARNING: Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle, the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Net Weight on Bag and/or Invoice

revision 12/20/16