



CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 654321 DATED 01/01/16



**JAMES SMITH GOAT FEED
MEDICATED**

TYPE C FEED CONTAINING RUMENSIN™(MONENSIN)

FORMULA #: SMITHGOATJan17

STATEMENT OF PURPOSE

For prevention of coccidiosis caused by Eimeria crandallis, Eimeria christenseni and Eimeria ninakohlyakimovae in goats maintained in confinement.

ACTIVE DRUG INGREDIENT

Monensin	20.00 grams/ton
	10.00 mg/pound

USE DIRECTIONS

Feed this complete feed (20 g/ton) continuously to goats as the sole ration.
Feed only to goats maintained in confinement.

PRECAUTIONARY STATEMENTS

CAUTION: Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

WARNING: Do not feed to lactating goats.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 5555

Net Weight on Bag and/or Invoice

RUMENSIN™ is a trademark of Eli Lilly & Company

Revision 8/8/2008



CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 654321 DATED 01/01/16



**JAMES SMITH GOAT FEED
MEDICATED**

TYPE C FEED CONTAINING RUMENSIN™(MONENSIN)

FORMULA #: SMITHGOATJan17

STATEMENT OF PURPOSE

For prevention of coccidiosis caused by Eimeria crandallis, Eimeria christenseni and Eimeria ninakohlyakimovae in goats maintained in confinement.

ACTIVE DRUG INGREDIENT

Monensin	20.00 grams/ton
	10.00 mg/pound

USE DIRECTIONS

Feed this complete feed (20 g/ton) continuously to goats as the sole ration.
Feed only to goats maintained in confinement.

PRECAUTIONARY STATEMENTS

CAUTION: Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

WARNING: Do not feed to lactating goats.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 5555

Net Weight on Bag and/or Invoice

RUMENSIN™ is a trademark of Eli Lilly & Company

Revision 8/8/2008