

|   |  |   |
|---|--|---|
|   | <b>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT</b><br>FOR INVOICE 20101 DATED 01/01/17 |  |
| <b>SMITH SWINE FARM SWINE FEED</b><br><b>MEDICATED</b><br>Complete Feed Containing Tiamulin (Denagard™) and Chlortetracycline   |  |   |
| <b>CAUTION:</b> Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.<br><b>VFD Expiration Date: Monday, January 30, 2017</b>   |  |   |
| FORMULA # SSF Denagard CTC Jan 17   |  |   |
| <b>STATEMENT OF PURPOSE</b>   |  |   |
| Treatment of swine bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline, and control of swine dysentery associated with <i>Brachyspria</i> (formerly <i>Serpulina</i> or <i>Treponema</i> ) <i>hyodysenteriae</i> susceptible to Tiamulin.  |  |   |
| <b>ACTIVE DRUG INGREDIENTS</b>  |  |   |
| Tiamulin  | 35.00 grams/ton  |   |
|   | 17.50 mg/pound   |   |
| Chlortetracycline   | 400.00 grams/ton   |   |
|   | 200.00 mg/pound  |   |
| <b>USE DIRECTIONS</b>   |  |   |
| Feed continuously as sole ration for 14 days as only source of chlortetracycline.   |  |   |
| <b>NOTICE: This feed may not be fed after</b> <span style="float: right;"><b>Monday, January 30, 2017</b></span>  |  |   |
| <b>PRECAUTIONARY STATEMENTS</b>   |  |   |
| <b>WARNING:</b> Withdraw 2 days before slaughter.<br><b>CAUTION:</b> Do not feed undiluted. Do not feed to animals other than swine. The effects of Tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin, and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.<br><b>Limitations:</b> Feed continuously as sole ration for 14 days. Feed approximately 400 g/T of chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. Drug combination approval only for ChlorMax, Aureomycin, or Pennchlor. |  |   |
| MANUFACTURED BY:  |  |   |
| BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555  |  |   |
| Net Weight on Bag and/or Invoice  |  |   |
| revision 12/1/2016  |  |   |

|   |  |  |
|---|--|--|
|   | <b>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT</b><br>FOR INVOICE 20101 DATED 01/01/17 |  |
| <b>SMITH SWINE FARM SWINE FEED</b><br><b>MEDICATED</b><br>Complete Feed Containing Tiamulin (Denagard™) and Chlortetracycline   |  |  |
| <b>CAUTION:</b> Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.<br><b>VFD Expiration Date: Monday, January 30, 2017</b>   |  |  |
| FORMULA # SSF Denagard CTC Jan 17   |  |  |
| <b>STATEMENT OF PURPOSE</b>   |  |  |
| Treatment of swine bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline, and control of swine dysentery associated with <i>Brachyspria</i> (formerly <i>Serpulina</i> or <i>Treponema</i> ) <i>hyodysenteriae</i> susceptible to Tiamulin.  |  |  |
| <b>ACTIVE DRUG INGREDIENTS</b>  |  |  |
| Tiamulin  | 35.00 grams/ton  |  |
|   | 17.50 mg/pound   |  |
| Chlortetracycline   | 400.00 grams/ton   |  |
|   | 200.00 mg/pound  |  |
| <b>USE DIRECTIONS</b>   |  |  |
| Feed continuously as sole ration for 14 days as only source of chlortetracycline.   |  |  |
| <b>NOTICE: This feed may not be fed after</b> <span style="float: right;"><b>Monday, January 30, 2017</b></span>  |  |  |
| <b>PRECAUTIONARY STATEMENTS</b>   |  |  |
| <b>WARNING:</b> Withdraw 2 days before slaughter.<br><b>CAUTION:</b> Do not feed undiluted. Do not feed to animals other than swine. The effects of Tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin, and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.<br><b>Limitations:</b> Feed continuously as sole ration for 14 days. Feed approximately 400 g/T of chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. Drug combination approval only for ChlorMax, Aureomycin, or Pennchlor. |  |  |
| MANUFACTURED BY:  |  |  |
| BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555  |  |  |
| Net Weight on Bag and/or Invoice  |  |  |
| revision 12/1/2016  |  |  |