



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WISCONSIN

Bravo Ultrex®

EPA Reg. No. 50534-201-100

EPA SLN No. WI-130009

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2017

For Control of Diseases on Long-Season Potatoes

**KEEP OUT OF REACH OF CHILDREN
DANGER/PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA- registered label.

This label must be in the possession of the user at the time of pesticide application.

It is a violation of Federal law to harm an endangered species. Use of this product may be in counties which have endangered and threatened species present; therefore, applicators should evaluate the areas to be treated and take necessary precautions to avoid harming endangered species or their habitat/environment. Applicators may refer to http://datcp.wi.gov/Farms/Endangered_Species/ for help in identifying the species and their habitats. Growers who are uncertain of the presence of a protected species should contact the Wisconsin Department of Agriculture, Trade and Consumer Protection at 608-224-4547.

Crop	Diseases (Pathogens)	Lb Product/A (lb ai/A)	Application Directions
Potato (Long-season varieties)	Late Blight (<i>Phytophthora infestans</i>)	Low rate: 0.7 (0.6)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days). Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur: <ul style="list-style-type: none"> Vines close between rows Late blight forecasting measures 18 disease severity values (DSV) The crop reaches 300 P-days Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe. Bravo Ultrex may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Ultrex label.
	Early Blight (<i>Alternaria solani</i>)	-then-	
	Botrytis vine rot (<i>Botrytis cinerea</i>)	Higher rates: 0.9 to 1.36 (0.75 to 1.1)	
	Black dot (<i>Colletotrichum coccodes</i>)		

Specific Use Restrictions: Do not apply more than 19.35 pounds of Bravo Ultrex (16 lb ai) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.

Bravo Ultrex® is a trademark of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300.
Greensboro, NC 27419-8300

Label Code: WI0201040AA0513