

# FIFRA 24(c) Special Local Need Label (SLN)

For Distribution and Use Only in Wisconsin

## Preplant Application to Control Soil Insects Infesting Ginseng

### Lorsban<sup>®</sup> 15G

EPA Reg. No. 62719-34

SLN WI-130006

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2016.

#### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban<sup>®</sup> 15G insecticide before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- Use of Lorsban 15G according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban 15G.

Apply Lorsban 15G only once per season. This application may occur as either a preplant treatment at an application rate of 13.5 lb (2 lb ai chlorpyrifos) per acre per application by 1) evenly broadcasting granules over the soil surface within 14 days of forming the plant beds and seeding the crop, followed by soil incorporation to a depth of 4 to 6 inches; 2) a bed-directed application immediately prior to seeding and mulching (but following bed formation); or 3) a post-emergence, bed-directed treatment by evenly distributing granules over the crop canopy followed by rainfall or an overhead watering applied at an application volume of 100 gallons per acre.

Target Pests	Lorsban 15G (lb/acre)
cutworms	13.5
grubs	
rootworms	
wireworms	

#### Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 365 days before harvest.
- Do not make more than one application of Lorsban 15G per crop.
- Maximum single application rate is 2 lb ai chlorpyrifos per acre.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry (coveralls, long-sleeved shirt, long pants, shoes and socks, chemical-resistant gloves, respirator) is worn.
- Individual workers applying Lorsban 15G to ginseng gardens may not treat more than 7 acres per day.
- **Chemigation:** Do not apply through any type of irrigation equipment.

It is a violation of federal law to harm an endangered species. Ginseng is grown in counties which have endangered and threatened species present; therefore, growers should evaluate their fields and take necessary precautions to avoid harming endangered species or their habitat/environment. Growers who are uncertain of the presence of a protected species should contact the Wisconsin Department of Agriculture, Trade and Consumer Protection at 608-224-4547.

**Environmental Hazards**

This pesticide is toxic to fish, aquatic invertebrates, small mammals and birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

® Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R061-018

Accepted: \_\_/\_\_/\_\_

Replaces: R061-016