



*Updated 3/30/22*

## **HPAI in Wisconsin: Frequently Asked Questions**

The Wisconsin Department of Agriculture, Trade and Consumer Protection (DATCP) will continue to update the following questions, in partnership with federal, state, and local partners.

### **How many facilities have tested positive for HPAI in Wisconsin?**

DATCP [announced](#) a confirmed case of highly pathogenic avian influenza (HPAI) in a commercial poultry flock in Jefferson County. Samples were tested by the Wisconsin Veterinary Diagnostic Laboratory and confirmed by the National Veterinary Services Laboratories in Ames, Iowa. At this time, the only confirmed positive in Wisconsin is the affected premises in Jefferson County. For regular updates see DATCP's [HPAI in Wisconsin](#) page.

### **Has HPAI been detected in the wild bird population?**

HPAI has been detected in the wild bird population and birds can carry the disease to new areas when migrating, potentially exposing domestic poultry to the virus. Additional avian influenza detections may occur as wild birds migrate.

USDA APHIS is conducting [wild bird surveillance](#), which serves as an early warning system for the introduction and distribution of avian influenza. The Wisconsin Department of Natural Resources (DNR) has also implemented enhanced wild bird surveillance in Wisconsin. To date, no wild birds in Wisconsin have tested positive.

### **How are wild bird populations being monitored?**

The Wisconsin Department of Natural Resources (DNR) is collaborating with federal partners by enhancing surveillance efforts. In addition to the normal morbidity and mortality investigations of five or more sick or dead birds of any species, the DNR is monitoring for possible incidents of EA H5 HPAI by focusing increased efforts on species that have been shown to be more likely to carry or be susceptible to disease from this strain.

Although there are no clinical signs specific to this HPAI strain, some affected birds may show neurologic signs. The public is asked to report any waterfowl, waterbird, raptor (especially bald eagles) or avian scavenger (crows, ravens, gulls) showing signs such as swimming or walking in circles, tremors, or holding the neck and head in an unusual position. Reports should be made to DNR Wildlife Hotline by email to [DNRWildlifeSwitchboard@wisconsin.gov](mailto:DNRWildlifeSwitchboard@wisconsin.gov) or by phone at (608) 267-0866. Leave a message with the number of animals; the species (e.g., Canada goose or mallard); if they were sick or dead; the specific location where you saw them, including the county; and the contact details of the person reporting so that staff can return a call. [More information is available on the DNR website here.](#)

## **How does avian influenza spread?**

Birds act as hosts to influenza viruses by carrying the virus in their intestines and shedding it in bodily fluids, such as saliva, nasal secretions and feces. Other birds become infected when they come in contact with these fluids. Humans can become infected through contact with infected poultry or contaminated fluids. Avian flu can spread from birds to people and cause serious illness and death.

Avian influenza has not mutated to a point where it could easily spread from person-to-person. It's difficult to predict if — or when — that might happen, or if it will result in an influenza pandemic. People who have gotten sick with avian influenza have been in direct contact with infected birds.

According to the CDC, HPAI does not present an immediate public health concern and no human cases of avian influenza have been found in the US. For more information, see the Wisconsin Department of Health Services (DHS) [avian influenza webpage](#).

## **Will HPAI enter the food system?**

Birds and eggs from the flock will not enter the food system. HPAI does not pose a food safety risk; properly handling and cooking poultry and eggs to an internal temperature of 165°F kills the virus.

## **What happens to the birds after depopulation?**

The birds will be composted; composting is the most efficient and environmentally responsible method for disposal of bird carcasses. Compost piles are constructed in a manner that includes a thick cover of compost carbon, such as wood chips, on top to eliminate the possibility of wild birds landing on the pile and carrying virus to new locations.

Compost piles are regularly inspected by staff from the farm, DNR, and DATCP for any evidence of leakage, odor, and disruption. The full composting process takes approximately 30 days from final construction and involves daily monitoring to ensure the compost is reaching the optimum temperature to deactivate the virus as quickly as possible. Windrows will reach temperatures exceeding 130 degrees for at least 72 hours and is sufficient to inactivate many pathogens including avian influenza.

After the composting process is complete and the compost has been released by DATCP, the compost will be removed from the premises at the discretion of the land owner.

DATCP has resources on composting, including resources from the University of Minnesota, Iowa State University, and Cornell University, on the [Livestock Carcass Disposal webpage](#).

## **How has the current compost site been chosen?**

The Wisconsin Department of Natural Resources (DNR), USDA, and DATCP evaluated several sites in the Jefferson County area relative to USDA animal mortality composting guidelines. According to the DNR, the preferred depth to groundwater should exceed 24 inches to seasonal highwater tables and on-site soil depths should exceed 36 inches to bedrock. The egg farm sits in

an area that does not meet this criteria. However, the composting site is owned by the farm and meets both the locational and size criteria for a compost site.

Additionally, the compost site should be:

- 200 feet from a water supply well used for drinking;
- 200 feet from water bodies, including ponds, lakes, streams and rivers;
- 200 feet from a nearby residence (not owned by the premises);
- 50 feet from a drainage swale that leads to a water body; and
- 25 feet from a drainage swale that does not lead to a water body.

DATCP will continue to update information specific to the compost site as it becomes available. To stay updated, [sign up to receive email updates](#) about HPAI in Wisconsin.

### **Why was composting selected as the disposal method instead of another disposal method?**

Composting is the safest and most effective way to handle large volumes of carcasses. The composting process deactivates the virus while minimizing odor and generation of liquids into the ground. Although some materials will go to landfills, landfills can generally only accept a limited amount of carcass material.

Rendering, burial, open-air burning and landfilling have been means of disposal, but are becoming less and less practical. Burial and open-air burning create biosecurity hazards and threats to water and air quality. To mitigate the spread of highly pathogenic avian influenza (HPAI) and safely dispose of the carcasses as soon as possible and due to the proximity and capability of incinerators in Wisconsin, incineration was not an option. For a large number of carcasses, such as the Jefferson County premises infected with HPAI, mobile incineration requires additional fuel and many units to be deployed.

### **What happens on a compost site?**

Material will be laid out in windrows, which are long rows of composting material. Each windrow will include a mixture of carbon (mostly woody/mulch materials) under, around, and on top of the carcasses to create the appropriate environment for effective composting. Encasing and topping the carcasses with at least one foot of the carbon layer encourages the higher temperatures that speed decomposition, absorbs odors, and provides additional protection against water penetrating to the area of the windrow where the chickens have been placed. For more information, see DATCP's [Livestock Carcass Disposal webpage](#).

### **Why will depopulated birds be moved to this site instead of being composted on the farm?**

Some have already been composted on-site within an existing building, the only appropriate area on the farm property. Lands surrounding the farm do not have a suitable depth to groundwater for on-site composting.

### **Why was this site selected for composting depopulated birds?**

The site is owned by the farm, has sufficient area, and meets the locational criteria for a compost site.

## **What is the risk of getting bird flu (avian influenza) from the groundwater?**

The risk of being infectedd with HPAI from groundwater is extremely low. There has never been a documented outbreak of HPAI from infected groundwater, and there has been no evidence of live viruses capable of infecting humans or animals.

Composting will rapidly deactivate HPAI and other pathogens when target temperatures are reached within the compost piles. Additionally, safety standards for this composting site were designed to prevent contamination of groundwater with HPAI and other byproducts of the composting process. These safety measures include proper construction of compost windrows to minimize the generation of liquids, and ensuring adequate distances are maintained from the compost site to well-water supplies, residences, and other bodies of water.

Resources:

- [EPA: Pandemic influenza fact sheet for water](#)
- [CDC: Prevention and transmission of bird flu](#)

## **How can I make sure my water is safe to drink?**

If your water comes from a private well, you should [test your water](#) regularly (at least once a year) for contaminants like bacteria, nitrates, and arsenic. If you suspect there is [a problem with your water](#), or are concerned about water contamination, you shouldn't wait to test your water. Although routine testing will not detect influenza viruses, the presence of bacteria would be an indicator of contamination and should be followed up with appropriate action to treat or disinfect your well. Annual testing of your well allows you to know your baseline levels so that any changes to your water quality can be recognized. You can find a list of certified labs that perform well water testing on the [DNR website](#).

Resources:

- [DNR: Information for private well owners](#)
- [DHS: Testing recommendations for private well owners](#)

## **What are the operational hours of the compost site?**

Working hours at the compost site will be 7 a.m. to 7 p.m.

## **How is HPAI being monitored in affected area?**

The affected premises is not allowed to move poultry or poultry products, and birds on the property will be depopulated to prevent spread of the disease. Birds from the flock will not enter the food system. Poultry premises within 10 kilometers of the farm are being monitored for the virus and are restricted from moving poultry and poultry products. Additionally, DATCP issued a special order banning poultry from any movement to, or participation in, shows, exhibitions, and swap meets held in Jefferson County. The special order will remain in effect through May 31, 2022.

## **How can I protect my poultry?**

DATCP continues to encourage everyone in Wisconsin working with or around poultry to practice enhanced biosecurity measures to protect their flocks, including:

- Sanitizing equipment and clothing used around your flock.

- Protecting your flock from contact with wild birds.
- Restricting access to your property and keeping your birds away from other birds.

For more information on biosecurity, see the biosecurity section of the DATCP's [HPAI in Wisconsin page](#).

### **What are the clinical signs of HPAI?**

- Sudden death, possibly without clinical signs
- Decrease in egg production or soft-shelled/misshapen eggs
- Nasal discharge, coughing, sneezing
- Swelling of the head, eyelids, comb, wattles, and hocks
- Lack of energy and appetite
- Difficulty breathing
- Stumbling, falling down
- Diarrhea

### **What should I do if I observe HPAI symptoms in my poultry?**

To report increased mortality or signs of illness among domestic birds, contact DATCP at (608) 224-4872 (business hours) or (800) 943-0003 (after hours and weekends).

### **What should I do if I find a sick or dead wild bird?**

You can [report sick or dead wildlife](#) (including birds) to the Wisconsin Department of Natural Resources (DNR). Depending on the type of bird, DNR may arrange to pick up the bird or instruct you to discard it. As a general precaution, avoid direct contact with wild birds whenever possible. Avoid touching any wild birds showing [signs of HPAI](#), and do not touch dead birds or wildlife with your bare hands. If you must touch a dead bird, wear gloves or use a plastic bag to put it in the garbage. Wash your hands with soap and water, and throw away any gloves after disposing of dead birds or wildlife.

### **More information**

DATCP has information and resources available on [datcp.wi.gov](http://datcp.wi.gov). With any additional questions, please contact [DATCPmediainquiries@wi.gov](mailto:DATCPmediainquiries@wi.gov). To receive daily updates, [sign up to receive email updates](#) about HPAI in Wisconsin.

DATCP will continue to update this document, in coordination with state and federal partners, as possible. As it is updated, DATCP will post it to the [HPAI in Wisconsin webpage](#).