

2021 Field-Edge Groundwater Monitoring Program

ANNUAL REPORT



Wisconsin Department of Agriculture, Trade and Consumer Protection
Agricultural Resource Management Division
Environmental Quality Unit
03-29-2023

Table of Contents

| | |
|---|-----------|
| List of Figures | 0 |
| List of Tables | 0 |
| Introduction | 1 |
| Purpose of Field-Edge Groundwater Monitoring | 1 |
| Program Approach | 2 |
| Program Assets and Infrastructure | 3 |
| 1985-1989 Original Monitoring Wells and Piezometers | 4 |
| 2005 Monitoring Program Expansion | 5 |
| 2010 University Wisconsin-Oshkosh Wells | 5 |
| 2011 Monitoring Program Expansion | 5 |
| 2017 Monitoring Program Expansion | 5 |
| 2021 Monitoring Program Expansion/Abandonment | 5 |
| 2021 Results | 6 |
| Grower Responses | 7 |
| Water Level Measurements | 8 |
| Pesticide Detection Frequency | 13 |
| Comparison to Standards | 14 |
| Other Notable Observations | 16 |
| Glyphosate: | 16 |
| Neonicotinoids: | 16 |
| Alachlor: | 17 |
| Atrazine: | 17 |
| Nitrogen: | 18 |
| 2022 Program Goals and Objectives | 22 |
| Acknowledgments | 23 |
| References | 23 |
| Appendix A | 24 |
| Acronyms | 24 |
| Definitions | 25 |
| APPENDIX B | 26 |

List of Figures

| | |
|--|----|
| Figure 1: 2021 Monitoring Well Sites _____ | 4 |
| Figure 2: Accumulated Precipitation from the Wisconsin Monthly Climate Watch Archive _____ | 8 |
| Figure 3: Monthly Precipitation Totals for Sampling-Site Counties from the NOAA Monthly Climate Watch Archive _____ | 9 |
| Figure 4: Wisconsin Monthly Precipitation Departures (from 1991-2020 Average) for 2021 _____ | 10 |
| Figure 5: Wisconsin Accumulated Precipitation (in): Departure from 1991-2020 Average _____ | 11 |
| Figure 6: Historic Water Table Level Data for a Field-Edge Monitoring Station AD2 in Adams County _____ | 12 |
| Figure 7: Historic Water Table Level Data for a Field-Edge Monitoring Station DU1 in Dunn County _____ | 12 |
| Figure 8: Historic Water Table Level Data for a Field-Edge Monitoring Station IW1 in Iowa County _____ | 13 |
| Figure 9: Percentage of 2021 Samples with Detectable Pesticide Concentrations (Includes all analytes detected in 20% or more of all samples collected) _____ | 14 |
| Figure 10: 2021 Atrazine TCR Concentrations relative to Groundwater Sample Well Depth _____ | 18 |
| Figure 11: Nitrogen as Nitrate plus Nitrite Results Distribution in Groundwater Samples from All-Wells _____ | 19 |
| Figure 12: 2021 Nitrogen as Nitrate plus Nitrite Concentrations relative to Groundwater Sample Well Depth _____ | 20 |
| Figure 13: 2021 Nitrogen as Nitrate plus Nitrite Concentrations Variability from Spring to Fall at Individual Wells _____ | 21 |
| Figure 14: 2021 Nitrogen Concentrations Variability by Depth from Spring to Fall of Individual Wells _____ | 22 |

List of Tables

| | |
|--|----|
| Table 1: Crops Grown and Nitrogen Applied on Fields Adjacent to Field-Edge Stations _____ | 7 |
| Table 2: Detected Parent Compounds that have No Wisc. Admin. Code ch. NR 140 Standard or WDHS Drinking Water Health Advisory Levels _____ | 15 |
| Table 3: Average Nitrogen as Nitrate plus Nitrite Concentration over Previous Years. _____ | 19 |
| Table B 1: Field-Edge Groundwater Monitoring Program - Monitoring Wells and Piezometers Construction Specifications _____ | 26 |
| Table B 2: 2021 Sample Analytes, Applicable Wis. Admin. Code ch. NR 140 PALs & ESs, Drinking Water Health Advisories, and Reporting Limits _____ | 28 |
| Table B 3: Field-Edge Groundwater Monitoring Program - 2021 Groundwater Analytical Results _____ | 29 |
| Table B 4: Field-Edge Groundwater Monitoring Program - 2021 Land Pesticide/Nitrogen- and Irrigation-Use (as Provided by Growers) _____ | 33 |
| Table B 5: Field-Edge Groundwater Monitoring Program - 2021 Imidacloprid Concentrations in Groundwater Samples _____ | 38 |
| Table B 6: Field-Edge Groundwater Monitoring Program - 2021 Alachlor ESA Concentrations in Groundwater Samples _____ | 40 |
| Table B 7: Field-Edge Groundwater Monitoring Program - 2021 Atrazine and Metabolite Concentrations in Groundwater Samples _____ | 42 |
| Table B 8: Field-Edge Groundwater Monitoring Program - 2021 Nitrogen-Nitrate/Nitrite Concentrations in Groundwater Samples _____ | 44 |

Introduction

In 2021, the Wisconsin Department of Agriculture, Trade and Consumer Protection's (DATCP) Agrichemical Management (ACM) Bureau continued the Field-Edge Groundwater Monitoring Program to document the effect continual pesticide use is having on groundwater quality. Groundwater monitoring was performed by DATCP staff across a network of 62 monitoring wells and piezometers at 24 established locations. At each location, depth to groundwater is measured and groundwater samples are collected in the spring and fall to identify pesticide concentrations and evaluate seasonal variations. Collected samples are submitted to DATCP's Bureau of Laboratory Services (BLS) for chemical analysis. This report has been prepared to document 2021 Program activities and includes a summary of groundwater level measurements and analytical data results. Recommendations for the 2022 Field-Edge Groundwater Monitoring Program plan, based on historic trend results, are also presented in this report.

A compilation of acronyms and definitions used throughout this document is provided in [Appendix A - Acronyms and Definitions](#).

Purpose of Field-Edge Groundwater Monitoring

It is estimated that agriculture contributes \$104.8 billion annually to Wisconsin's economy (Wisconsin Department of Agriculture, Trade and Consumer Protection, 2023a). Growers in Wisconsin use several million pounds of pesticides and tons of fertilizers annually to grow a wide variety of crops. DATCP's Field-Edge Groundwater Monitoring Program is one form of monitoring the agency performs to meet its statutory obligation to protect groundwater quality. Wisconsin's groundwater law, Chapter 160, Wis. Stats., requires agencies to sample and monitor groundwater for substances related to facilities, activities, and practices under their jurisdiction, that have a reasonable probability of entering the groundwater resources of the state, and to determine whether preventive action limits (PAL) or enforcement standards (ES) have been exceeded at points of standard application. The statute further specifies that agencies should develop monitoring plans that include provisions for conducting four types of monitoring (Wis. Stats., Ch. §160.05 and §160.27):

1. Problem assessment monitoring, to detect substances in the groundwater and to assess the significance of the concentrations of the detected substances;
2. Regulatory monitoring, to determine if preventive action limits or enforcement standards are attained or exceeded and to obtain information necessary for the implementation of responses with respect to specific sites;
3. At-risk monitoring, to define and sample at-risk potable wells in areas where substances are detected in the groundwater or where preventive action limits or enforcement standards are attained or exceeded; and
4. Management practice monitoring, to assure practices are within compliance regulations.

The purpose of the Field-Edge Groundwater Monitoring Program (Program) is to evaluate agricultural practices and agrichemical uses on groundwater quality (problem assessment and regulatory monitoring). Depth to groundwater measurements and groundwater sample results are used to measure affects from agrichemical practices and use within and adjacent to agricultural fields. Affects to groundwater quality from agrichemical use is dependent on conditions at each location. Results are used to measure both localized and regional affects to aquifers over time at each field-edge sampling site. Historic and current goals of the Program include the following:

- Provide an early warning system to detect new agrichemical compounds in groundwater before widespread contamination can occur in underlying aquifers.
- Identify and measure pesticide concentrations that may have a potential to migrate to groundwater and exceed groundwater quality standards.
- Identify which environmental conditions (i.e. depth to groundwater, soil type, and geologic setting) are most vulnerable to conditions from routine agrichemical use.
- Gather and compile data regarding the occurrence and persistence of pesticide and metabolites in groundwater that may affect drinking water wells so that health-based groundwater quality standards can be established.

- Study the dissipation of restricted use pesticides (i.e. atrazine) in groundwater after prohibition areas are established or use is restricted, and the dissipation of pesticides no longer in use (i.e. aldicarb).
- Gather and compile long-term data on nitrate contamination in groundwater and its relationship to application practices.
- Evaluate affects to groundwater quality from various land uses and related pesticide use (i.e. tree nurseries, infiltration basins, golf courses).

Program Approach

DATCP and the property owner typically have access agreements allowing DATCP to install and access wells for sample collection. Typically, a monitoring well nest consists of a shallow well intersecting the water table and adjacent deeper wells (piezometers) installed with well screens placed at deeper depths within the underlying aquifer. These well nests are installed at the edge of an agricultural field to measure potential affects from routine agrichemical use. Well locations were carefully selected to avoid interference from other potential sources (i.e. septic systems).

Over time, monitoring well nests have been installed within a variety of geologic settings, often in areas prone to groundwater contamination, such as areas with sandy soil, shallow depths to bedrock, or shallow groundwater. Nested well locations have two to five monitoring wells/piezometers. The shallowest well intersects the water table with piezometers installed at deeper intervals. [Table B 1](#) in [Appendix B](#) provides construction specifications for each well in the Program's groundwater monitoring well network. [Figure 1](#) depicts the Program's monitoring locations relative to State of Wisconsin and county boundaries.

Program data collection and documentation are completed in accordance with established protocols and guidance (Wisconsin Department of Agriculture, Trade and Consumer Protection, 2021; Wisconsin Department of Natural Resources, 1996). Depth to water measurements and sample collection procedures are designed to collect reliable data consistently and in an unbiased fashion to ensure that localized conditions and regional impacts to aquifers over time can be evaluated. Field sampling observations and water level measurements are recorded in field notebooks. The compiled field information, along with laboratory results, are retained in databases maintained by DATCP.

Standard operating procedures for groundwater sampling include the following:

- After unlocking the protective casing, remove the well cap to allow the water level to equilibrate with atmospheric pressure before measuring and recording the water level at each well.
- Each well is then properly purged to remove a minimum of four well casing volumes. Purging is performed either by using dedicated bailers and rope, peristaltic pumps (low flow) with dedicated tubing, or submersible electric pumps (i.e. whale or tornado pumps) with dedicated tubing. The volume of water removed is measured and recorded in the field logbook.
- Samples are then collected and placed in laboratory-provided containers using either sampling equipment dedicated to the well, or with equipment that is decontaminated prior to use.
- Samples are placed into coolers and held on ice while in transport to the laboratory.
- Water purged from the wells and any rinse water used for cleaning is discarded on the ground surface.
- Field information is recorded in logbooks and maintained by ACM staff.

Groundwater samples are collected using the same equipment used for purging. Samples are collected in one-liter amber glass bottles provided by BLS. (Fifty-millimeter plastic containers were used for select glyphosate sampling.) Bottles and containers are then placed in a cooler and held on ice along with a properly completed sample collection record and hand-delivered to BLS within 48 hours. During the 2021 Program, there were no issues with shipping or bottle breakage.


BLS performed all groundwater analytical testing using gas chromatography/mass spectroscopy (GC/MS/MS) and liquid chromatography/mass spectroscopy (LC/MS/MS) methods in accordance with ISO 17025 accreditation standards. All samples were tested for 106 pesticide analytes as well as nitrogen as nitrate plus nitrite. Pesticide analytes are listed in [Table B 2](#) of [Appendix B](#) along with corresponding reporting limits. A summary of the 2021 Program analytical data results is listed in [Table B 3](#) of [Appendix B](#). Individual monitoring well or piezometer analytical reports are available upon request.

DATCP provides annual Program findings documentation for each site to the respective property owner or grower. The summary letters provide the year's water level data and analytical results, and includes a brief discussion of data trends over time. As part of the letter, growers are asked to reply with information regarding crops grown, pesticide use, and the amount of nitrogen applied to the fields near monitoring wells.

Program Assets and Infrastructure

The groundwater-monitoring network for the 2021 Field-Edge Monitoring Program included 62 groundwater monitoring wells (31 water table observation wells and 42 piezometers) at 24 locations/stations around the state. [Table B 1](#) in [Appendix B](#) lists well construction specifications associated with these Program assets. [Figure 1](#) depicts the Program's monitoring sites relative to State of Wisconsin and county boundaries. Construction logs and well development forms (and abandonment forms) associated with the groundwater monitoring wells and piezometers are available upon request. The following is a summary of the Program's well installation history.

Figure 1: 2021 Monitoring Well Sites



1985-1989 ORIGINAL MONITORING WELLS AND PIEZOMETERS

The DATCP Field-Edge Groundwater Monitoring Program began in 1985. Initially, arrangements with growers and landowners at 50 sites were established in areas highly susceptible to groundwater contamination (i.e. coarse soil over sand, shallow to groundwater and/or irrigated agricultural areas). Groundwater monitoring nests with three to four wells were installed at each site. Nested wells were constructed with well screens placed at various depths in the underlying aquifer. These wells were constructed adjacent to agricultural fields in the Central Sands region, Lower Wisconsin River Valley, and at other sandy soil areas throughout the state. The original Field-Edge Study was designed to collect groundwater samples from the uppermost shallow aquifer. Samples were tested for a limited number of agrichemicals and fertilizer to evaluate potential impacts to shallow groundwater from routine agricultural practices performed at nearby fields.

Data from the Program's initial years led to the establishment of statewide pesticide management plans for both atrazine and aldicarb. Over the years, many of the wells installed for the initial study have been abandoned due to changes in land ownership, urban encroachment, or damage. Of the original 50 sites, monitoring wells remain at 16 sites and were included in the 2021 monitoring Program.

2005 MONITORING PROGRAM EXPANSION

In the fall of 2005, DATCP expanded its groundwater monitoring network with funding from a United States Environmental Protection Agency (US EPA) grant. New monitoring wells and piezometers were constructed at six sites based on local agricultural practices and susceptible to groundwater contamination (i.e. shallow groundwater with permeable subsurface soil units). Each of the six sites selected for Program expansion were used for a prior groundwater monitoring study (Evaluation of Renewed Use of Atrazine in Atrazine Prohibition Areas), completed by DATCP in 2005. That study (also known as the Atrazine Reuse Study) was performed to gather information to evaluate the potential to repeal atrazine prohibition areas.

The groundwater flow direction was determined as part of the Atrazine Reuse Study. Using that information, two monitoring wells were installed hydraulically down gradient and adjacent to agricultural fields at the six new sites. All six of these sites still were included in the 2021 monitoring Program.

2010 UNIVERSITY WISCONSIN-OSHKOSH WELLS

In the spring of 2010, DATCP became aware of a forthcoming study by a University of Wisconsin-Oshkosh graduate student and the Wisconsin Geological and Natural History Survey (WGNHS). The study included installation of shallow bedrock monitoring wells at the edge of agricultural fields in a karst geological setting. It used monitoring wells at sites in Brown, Calumet, Kewaunee, and Manitowoc counties. Bedrock fractures at each well were identified by the study team. Groundwater samples were collected by the study team and DATCP, and tested annually as part of this Program between 2010 and 2014. The study was completed and all monitoring wells were subsequently abandoned in 2014.

2011 MONITORING PROGRAM EXPANSION

In the summer and fall of 2011, DATCP expanded its groundwater monitoring network again with additional funding from an US EPA grant. Monitoring wells were constructed at two new stations in La Crosse and Trempealeau Counties. These wells were installed along an elevated terrace adjacent to the Mississippi River. Since the groundwater flow direction was known at each site (both locations were part of the Atrazine Reuse Study), DATCP installed two groundwater monitoring wells at each site at the hydraulically down gradient edge of each agricultural field. Wells at both sites remain and were included in the 2021 Program.

2017 MONITORING PROGRAM EXPANSION

In the summer and fall of 2017, DATCP further expanded the groundwater monitoring network with additional funding from a US EPA grant. Piezometers were constructed at three existing sites (two sites in Adams County and one in Portage County) and at one new site, the Hancock Agricultural Research Station (HARS). At each of these sites, two piezometers were installed near the existing groundwater monitoring nest with five-foot screens located at depths greater than 50 feet and 80 feet. The purpose was to evaluate groundwater quality relative to agrichemical uses at deeper aquifer intervals and compare data to shallower aquifer depths. A water table observations well (well screen placed to intersect the water table) was also constructed at HARS. The HARS site and nested wells at the Adams and Portage County sites remain and were included in the 2021 Program.

2021 MONITORING PROGRAM EXPANSION/ABANDONMENT

In the summer and fall of 2021, DATCP obtained additional funding from a US EPA grant again to expand the groundwater monitoring network. Eleven monitoring wells/piezometers were installed at six existing nested monitoring well sites. New wells were installed at sites in Adams County (AD2 and AD5), Dane County (DN1), Sauk County (SK6), Waushara County (WS7), and at two sites in Iowa County (IW1 and IW2). A monitoring well was also installed at the Dane County site to replace a well that was damaged beyond repair and subsequently abandoned in 2018. This shallow well was installed with a well screen intersecting the water table. Wells installed at the other five sites were constructed as piezometers with well screens placed 30 to

40 feet further in depth below the deepest existing piezometer screen already on-site in the well nest. These new piezometers were constructed with 5-foot long well screens. The purpose was to evaluate groundwater quality relative to agrichemical uses at deeper aquifer intervals and compare data across vertical aquifer horizons. All new wells were included in the 2021 fall sampling event.

Additionally, five wells at two monitoring locations were removed from the Program in 2021 in response to a change in property ownership. New owners for two Adams County sites (AD3 and AD4) did not want to continue to participate in the Program and requested removal of the wells. Two shallow water table observation monitoring wells and three piezometers were abandoned in December 2021.

2021 Results

A total of 126 water level measurements and 106 groundwater samples were collected as a part of DATCP's 2021 Field-Edge Groundwater Monitoring Program. All groundwater samples were submitted to BLS for chemical analysis. [Table B 3](#) in [Appendix B](#) summarizes 2021 Program analytical results and provides comparative risk values. The analytical data is compared to groundwater/drinking water standards to assess potential risk to human health and the environment. The risk values are sourced from the Wisc. Admin. Code ch. NR 140 for groundwater qualitative health standard limits and Wisconsin Department of Health Services (DHS) drinking water health advisories.

Key findings for 2021 include the following.

- One groundwater monitoring well (DN1) and seven piezometers (AD2, AD5, IW1, IW2, SK6 and WS7) were constructed at existing monitoring nest locations in 2021 and added to the Program. Groundwater samples for these locations were only collected during the fall sampling.
- Two monitoring nest locations (AD3 and AD4) were removed from the Program. Wells and piezometers were properly abandoned in December 2021. Groundwater samples were only collected during the spring sampling event from these two locations.
- Information regarding field use of pesticides and fertilizer was requested from growers for 23 sites, but only eight growers responded.
- Water level measurements show a slight decline in water table elevations in 2021 due to reduced precipitation compared to prior years. In 2021, according to National Oceanic and Atmospheric Administration (NOAA), the state received on average 3.9 inches of precipitation less than normal conditions. Above-average precipitation levels were recorded during the prior five years.
- Laboratory analysis include 106 pesticide analytes for the laboratory testing methods. During 2021, 33 pesticide analytes were detected in excess of reporting limits in numerous groundwater samples, which is similar to previous years.
- Pesticides detected in 2021 samples in excess of laboratory reporting limits include 13 herbicides, 13 herbicide metabolites, six insecticides, and one fungicide.
- It appears that pesticides were detected at slightly greater concentrations during the fall sampling event compared to spring results.
- Overall, analytical data collected at nested monitoring wells indicates that pesticide and nitrogen concentrations increase with depth. Greater concentrations at depth indicate that pesticides migrate vertically and laterally within the underlying aquifers. This trend is consistent with prior years' findings. However, the greatest pesticide and nitrogen concentrations (aside of atrazine and its metabolites) were not observed in the deepest wells installed in 2021. New monitoring wells were constructed in 2021 with screens at deeper depths.
- Metolachlor ethanesulfonic acid (ESA) was detected in excess of laboratory reporting limits in 98% of all samples collected, and was the most frequently detected pesticide in 2021. Additionally, ESA was detected at each groundwater monitoring site. This is consistent with prior years' findings.
- Clothianidin was the second most frequently detected compound. It was detected in excess of laboratory reporting limits in 75% of the samples collected, and at 20 of the 24 groundwater monitoring sites. These observations are consistent with findings from prior years.
- Alachlor ESA was the third most frequently detected compound. It was detected in excess of laboratory reporting limits in 65% of the samples collected, a 10% decrease in detection rate compared to the

previous year. However, the number of sites where it was detected (19 sites) is consistent with the prior year’s findings.

- Atrazine concentrations or one of its breakdown products (de-ethyl atrazine, de-isopropyl atrazine and diamino atrazine) was detected in excess of laboratory reporting limits in 46% of the samples collected. At each site with nested wells, results were evaluated by well depth. The greatest concentrations were detected in groundwater samples collected from the deepest piezometers.
- Neonicotinoid compounds clothianidin, imidacloprid and thiamethoxam were detected in excess of laboratory reporting limits in 75%, 50% and 45%, respectively, of the samples collected in 2021. The frequency of detection is similar to observations from the previous year.
- There were no Wisc. Admin. Code, ch. NR 140 ES exceedances of established groundwater quality health standards. (Note; only 31 of the 106 pesticides tested for have established groundwater quality health standard levels). However, there were exceedances of Wisc. Admin. Code, ch. NR 140 PAL for alachlor ESA, atrazine, de-ethyl atrazine, de-isopropyl atrazine, di-amino atrazine, and atrazine total chlorinated residuals (TCR).
- The Wisconsin Department of Health Services (WDHS) has also established drinking water quality advisories for several pesticides. Imidacloprid was detected at 14 out of 24 sites, with 12 of the 14 samples exceeding the WDHS drinking water health advisory level of 0.2 micrograms per liter (µg/L) or parts per billion (ppb).

GROWER RESPONSES

DATCP obtained limited information for 2021 regarding crops grown, pesticide use, and the amount of nitrogen applied to the fields adjacent to monitoring wells. A request for this information was included with each summary letter sent to nearby property owners and growers. Responses to the information request is voluntary. DATCP received replies from eight of the 23 sites. No information was requested from HARS for site WS7. [Table B 4](#) in [Appendix B](#) summarizes information provided by the growers along with available information from the previous four years. The following [Table 1](#) is a summary of crops grown adjacent to the monitoring well nests and nitrogen use data for 2021.

Table 1: Crops Grown and Nitrogen Applied on Fields Adjacent to Field-Edge Stations

| Crop | Number of Sites with Crops | Percent of Sites (reported) | Range of Nitrogen Applied (lbs / acre) |
|------------|----------------------------|-----------------------------|--|
| Corn | 4 | 50% | 133 - 518 |
| Potatoes | 1 | 12.5% | 282.9 |
| Snap Beans | 2 | 25% | 65-152.5 |
| Soybeans | 1 | 12.5% | 0 |

Irrigation systems are present at 19 of the 24 monitoring sites. Of the 19 sites with irrigation systems, seven sites provided water usage data for 2021. Growers reported that the range of irrigation water applied to the fields in 2021 ranged from 4.2 to 15.60 inches per acre.


Growers were also asked if they have state-approved Nutrient Management Plans for the adjacent fields. Of the eight respondents, only one indicated they have an approved plan.

A wide variety of pesticides used on fields adjacent to field-edge monitoring wells was reported by the growers. Metolachlor was the most widely used active ingredient pesticide followed by glyphosate. A total of 26 different active ingredients (pesticide compounds) were reported to be applied in 2021 to the fields. [Table B 4](#) in [Appendix B](#) identifies the complete list of pesticides used in 2021 as reported by the growers.

WATER LEVEL MEASUREMENTS

Depth to water level measurements are recorded for each well prior to collecting groundwater samples for laboratory analysis. Water level data is incorporated into a DATCP database for evaluation of historic trends. Water level data for 2021 was measured in the late spring (April and May), and fall (October and November). Overall, water level measurements for 2021 show declining trends, but remained at or above the average elevation compared to historic readings. Wisconsin averages 33.5 inches of precipitation annually. In 2021, the state of Wisconsin as a whole experienced below-average precipitation levels; 29.6 inches of rain (Wisconsin State Climatology Office, 2023). Shown in Figure 2 at a more localized level is the total accumulated precipitation, mapped across Wisconsin by the Wisconsin State Climatology Office. As indicated, there was an uneven distribution of rainfall across the state in 2021. The northern half of the state accumulated between 30 to 40 inches of precipitation while the southern half accumulated 15 to 25 inches of precipitation in 2021.

Figure 2: Accumulated Precipitation from the Wisconsin Monthly Climate Watch Archive




As recorded by NOAA, Figure 3 summarizes the 2021 total annual precipitation in the counties where Program groundwater monitoring stations are located. The various colors indicate the monthly precipitation data at each location.

It was reported in the NOAA Storm Events Database, from January through late February, that numerous heavy snow events occurred in northern Wisconsin while the lower portions of the state received minimal snowfall (NOAA - National Oceanic and Atmospheric Administration, 2023a). In late May, there were multiple heavy rain events from Crawford County to La Crosse County, cumulatively receiving 9 to 13 inches of rain in less than a week, resulting in flash floods. In late June through early July, multiple storms moved across western Wisconsin, with La Crosse to Eau Claire County impacted the most, resulting in a rapid accumulation


of 2 to 6 inches of rain, flash floods, and streambank erosion. In late July, a storm event in eastern Wisconsin from Fond du Lac to Milwaukee County resulted in urban and stream flooding. Although occasional storms were recorded, from April through December, it was a dry season for southern Wisconsin, where droughts were declared, and southeastern Wisconsin reported severe (D2) to extreme (D3) drought conditions, indicating potentially significant crop die-off according to the National Integrated Drought Information System (NOAA - National Oceanic and Atmospheric Administration, 2023b). The remainder of precipitation throughout the year, from August through December, primarily consisted of minor seasonal storm events.

Figure 3: Monthly Precipitation Totals for Sampling-Site Counties from the NOAA Monthly Climate Watch Archive



Monthly statewide precipitation departure from the historical normal was obtained from the Wisconsin State Climatology Office and is displayed on Figure 4 (Wisconsin State Climatology Office, 2023). During 2021, January, February, April, May, September, October, and November reflected a positive departure from normal, meaning that there was a decrease in precipitation compared to average. These range from -0.2 to -1.6 inches less than normal. Conversely, March, July, August, and December showed a positive departure from normal, meaning there was an increase in precipitation. These values ranged from 0.4 to 1.6 inches.


Figure 4: Wisconsin Monthly Precipitation Departures (from 1991-2020 Average) for 2021



Similarly, [Figure 5](#) depicts the departure from normal for the accumulated precipitation for 2021 data. Positive values, indicated in green, represent where total precipitation for the year was greater than average; negative departures, indicated by the yellow and orange colors, represent areas where total precipitation was lower than average. Notably, southern Wisconsin experienced 10 to 17 inches of total precipitation less than normal, shown in [Figure 5](#). According to NOAA’s *Annual 2021 National Climate Report*, Wisconsin accrued greater than a 3.90-inch deficit relative to normal conditions. This is the first negative precipitation total in the state of Wisconsin since 2012 (National Centers for Environmental Information (NCEI), 2021).

Figure 5: Wisconsin Accumulated Precipitation (in): Departure from 1991-2020 Average


January 01, 2021 to December 31, 2021



The following [Figures \(6 to 8\)](#) provide examples of measured water level fluctuations over time for three wells in the groundwater monitoring network. These three wells are at sites with infrastructure for irrigation. Growers responded to requests for information for the DU1 and IW1 sites; there was no irrigation water usage data provided for site AD2. Graphs showing water level measurement trends for all other wells in the groundwater monitoring network are available upon request.


2021 water level data for the Adams County station indicate a lowering water level relative to the immediate past ([Figure 6](#)). However, 2021 water levels are still greater than the average for the duration of the monitoring Program. There was an average level of precipitation in the area in 2021, recorded at 32 inches, versus above average levels in the past couple of years.

Figure 6: Historic Water Table Level Data for a Field-Edge Monitoring Station AD2 in Adams County



2021 water level data for a Dunn County station also indicated a continued decrease compared to the previous year (Figure 7). However, 2021 water levels are still greater than the average over the duration of the monitoring Program. As listed in the grower response, the adjacent property owners reportedly irrigated 15.6 inches of water on the corn crop in 2021. This volume is the greatest amount of water used for irrigation on the property over the past five years, which had ranged from 0.8 to 3.97 inches.


Figure 7: Historic Water Table Level Data for a Field-Edge Monitoring Station DU1 in Dunn County



2021 water level data for an Iowa County station indicates stable water table conditions, consistent with historical measurements (Figure 8). Because this site is near the Wisconsin River, it is likely influenced by river water levels. High water table conditions in the spring have been observed several times at this location over the course of the monitoring Program. The overall trend continues to indicate a stable trend over the past 20 years, which likely correlates to nearby river elevations. As listed in the grower response,

the adjacent property owners reportedly irrigated 9.4 inches of water on the corn crop in 2021. This is within the range of irrigation reported for this property in the past five years, which ranged from 5.7 to 21 inches of water.

Figure 8: Historic Water Table Level Data for a Field-Edge Monitoring Station IW1 in Iowa County




DATCP is planning to complete an additional evaluation of groundwater elevation data for each individual monitoring site as part of a detailed study. Historical water level monitoring data will be evaluated for each site and results will be documented in a separate report prepared for each site (*Historical Field-Edge Site Data Analysis*). This evaluation will include a comparison of water level trends to precipitation records. These reports are planned to be completed over a three-year period with the first group available in 2023.

PESTICIDE DETECTION FREQUENCY

Thirty-three of the 106 analytes tested in DATCP’s 2021 Field-Edge Groundwater Monitoring Program were detected in excess of laboratory reporting limits. The number of compounds detected in 2021 increased from 28 pesticides detected in the prior year. Clopyralid, dimethenamid OA, and prometon were detected for the first time in the field edge monitoring pProgram in 2021. A pesticide analyte was detected in every groundwater sample collected in the 2021 Field Edge Program, with exception of two samples collected from the monitoring well network located in Barron County. Pesticides detected in excess of laboratory reporting limits in 2021 samples include 13 herbicides, 13 herbicide metabolites, six insecticides, and one fungicide.

The most frequently detected pesticide compounds detected in 2021 are listed in [Figure 9](#). This includes all pesticide analytes detected at a concentration greater than the laboratory reporting limit at a frequency greater than 20%. The number of compounds detected at this rate is increased in 2021 compared to prior years.

Figure 9: Percentage of 2021 Samples with Detectable Pesticide Concentrations (Includes all analytes detected in 20% or more of all samples collected)



Notes: *Atrazine TCR is total chlorinated residues of atrazine, which includes the sum of atrazine plus its metabolites de-ethyl atrazine, de-isopropyl atrazine, and di-amino atrazine.*

Metolachlor ESA was the most frequently detected analyte in excessive of laboratory reporting limits. It is a breakdown product of metolachlor, which is an active ingredient in corn herbicides. Metolachlor ESA was detected at every site and in 98% of all samples collected.

Clothianidin was the second most frequently detected compound in 2021. It was detected in excess of laboratory reporting limits at 20 of the 24 sites and in 75% of the samples collected. This rate of detection represents a continuing increase of clothianidin detections since clothianidin testing began 14 years ago. In previous years, clothianidin detections were commonly observed at sites within the Central Sands Agricultural Region, but rarely observed elsewhere. Clothianidin is now widely detected at most field-edge monitoring well sites within agricultural-intense areas.

The third most frequently detected analyte for the 2021 Program was alachlor ESA. It was detected in excess of laboratory reporting limits at 19 of 24 sites and in 65% of the samples collected. This represents a 10% decrease in the number of alachlor ESA detections since 2020.

COMPARISON TO STANDARDS

The Wisconsin Department of Natural Resources (WDNR) sets groundwater quality standards in Wisc. Admin. Code ch. NR 140, which includes substances of public health concern based on recommendations from WDHS. These standards have two parts, the ES and the PAL. The ES is a level that, if exceeded, requires intervention from the appropriate authority. In the case of pesticides in drinking water, DATCP is required to intervene if levels exceed the ES. The PAL is a percentage of the ES: 10% of the ES for carcinogenic, mutagenic or teratogenic properties and 20% of the ES for all other substances. The intention of the PAL is

to act as a trigger for intervention by the appropriate authority before the pollutant becomes a risk to public health.

Pesticide concentrations identified during DATCP’s 2021 Program were compared to Wisc. Admin. Code ch. NR 140 Groundwater Quality standards. WDHS has also established drinking water quality advisories for 15 different pesticides. [Table B 3](#) in [Appendix B](#) lists the existing standards alongside the range of concentrations for the pesticide compounds detected in 2021 groundwater samples

No ES standards were exceeded in any samples collected in 2021. However, imidacloprid concentrations exceeded the WDHS drinking water health advisory of 0.2 µg/L in 12 groundwater samples collected from nested monitoring wells sites in Adams, Iowa, Sauk, and Waushara counties. These sites are located in the Lower Wisconsin River Valley and the Central Sands Agricultural Region. Imidacloprid concentrations ranged from 0.204 to 2.77 µg/L. No other WDHS drinking water health advisories were exceeded in 2021 samples.

As depicted in [Table B 3](#) in [Appendix B](#), concentrations of alachlor ESA, atrazine, de-ethyl atrazine, de-isopropyl atrazine, di-amino atrazine, and atrazine TCR (total chlorinated residues, which are the sum of atrazine plus its metabolites de-ethyl atrazine, de-isopropyl atrazine, and di-amino atrazine) were detected in excess of the Wisc. Admin. Code ch. NR 140 PAL standards. The locations of the wells with PAL exceedances and detected concentrations are fairly consistent with results from prior years.

[Table B 3](#) in [Appendix B](#) also includes results for pesticides and their metabolites with no established ES or PAL. 74 out of 106 pesticides compounds tested have no established standard. A review of all 2021 data indicates that 34 different pesticides compounds were detected in excess of laboratory reporting limits, and 16 of these 34 compounds have no Wisc. Admin. Code ch. NR 140 established standard. However, nine of the 16 compounds with no established standard have a WDHS drinking water health advisory (clothianidin, imidacloprid, sulfentrazone, thiamethoxam, chlorantraniliprole, flumetsulam, fomesafen, metalaxyl, and saflufenacil). Four of the 16 compounds with no established standards or WDHS advisories are metabolites for compounds with standards (alachlor, dimethenamid, or metribuzin). The remaining four detected compounds with no existing standard or WDHS advisory are bicycloprone, imazethapyr, clopyralid, and cyantraniliprole. [Table 2](#) includes a detection summary of these remaining four compounds that are not metabolites and have no standard or advisory.

Table 2: Detected Parent Compounds that have No Wisc. Admin. Code ch. NR 140 Standard or WDHS Drinking Water Health Advisory Levels

| Analyte | Sites with Detects (out of 24) | Number of Detects (out of 80) | % of Samples Detected | Concentration Range (in µg/L) |
|------------------|--------------------------------|-------------------------------|-----------------------|-------------------------------|
| Bicycloprone | 2 | 2 | 1.9% | 0.0736-0.0802 |
| Imazethapyr | 1 | 2 | 1.9% | 0.117-0.228 |
| Cyantraniliprole | 4 | 5 | 4.7% | 0.0515-0.224 |
| Clopyralid | 1 | 1 | 0.9% | 0.32 |

This is the first time clopyralid, prometon, and dimethenamid OA, have been detected in excess of laboratory reporting limits in any of the groundwater samples associated with the Program. Clopyralid is a broadleaf herbicide intended for thistles and clovers, prometon is a non-selective herbicide for non-crop areas such as paths or around buildings, while dimethenamid OA is an herbicide metabolite.

It is important to note that comparisons of detected pesticides and their metabolite concentrations to established groundwater quality standards and drinking water advisories are based on exposure to a single compound. These comparisons do not fully evaluate the risk to human health when two or more compounds are present. Currently, there are no calculations to predict potential risk when multiple compounds are present. Since the current approach does not account for potential cumulative risk, potential toxicity may be underestimated when two or more compounds are present.

OTHER NOTABLE OBSERVATIONS

Glyphosate:

According to USDA (United States Department of Agriculture) - National Agricultural Statistics Service, in 2020, glyphosate was the most widely used pesticide on Wisconsin fields planted with soybean and second most pesticide used on fields planted with corn (United States Department of Agriculture - National Agricultural Statistics Service, 2023). Until 2019, glyphosate and the metabolites were not included in the DATCP pesticide analysis. Because glyphosate has been widely used (and has been for many years prior), DATCP added limited testing for glyphosate and two of its metabolites, AMPA (aminomethylphosphonic acid) and glyphosate ammonium, to the 2019 testing Program.

For 2021, glyphosate sampling was limited to 14 samples collected in June and October from monitoring wells at seven different locations (DU1, DU2, JN1, JN3, LC2, SC1 and TR1). In addition to the full pesticides scan, these samples were also tested for glyphosate and its metabolites. Based on the crops grown or as reported by the growers in their Response Reports ([Table B 4 in Appendix B](#)), glyphosate would or could have been applied to these adjacent fields in 2021 and the previous years. No detections in excess of laboratory reporting limits for any of the glyphosate family of pesticides were reported in these groundwater samples collected in 2021.

Neonicotinoids:

Interest in the neonicotinoid class of insecticides has increased greatly in recent years due to concerns over possible effects on pollinators. DATCP began testing for these compounds in 2008 with thiamethoxam. BLS now analyzes for six neonicotinoid compounds. Three of these compounds - clothianidin, imidacloprid, and thiamethoxam (CIT) - were detected in field-edge groundwater samples collected in 2021. The other three neonicotinoid compounds - acetamiprid, dinotefuran, and thiacloprid - were not detected in excess of laboratory reporting limits in any groundwater samples. The presence of the three CIT compounds in groundwater is expected as these compounds are known to readily leach when applied to crops grown in sandy soils and are used in many insecticide products. CIT compounds are labeled for use on most crops grown in the state including corn, soybeans, potatoes, many other vegetables, fruit crops, and most small grains.

Historic field-edge monitoring results indicate that CIT compounds are becoming more prevalent in groundwater over time. CIT compounds were observed at more locations in 2021 compared to prior years. However, concentrations seem to be stable or slightly decreasing at areas with known impacts. Thiamethoxam and imidacloprid have been detected in field-edge samples since testing for neonicotinoid compounds began primarily at sites within the Central Sands Agricultural Region and Lower Wisconsin River Valley.

No Wis. Admin. Code ch. NR 140 ES or PAL groundwater quality standards have been established for the CIT compounds. However, DHS has identified drinking water health advisories for the CIT compounds.

Clothianidin and thiamethoxam were detected in 75% and 45%, respectively, of all 2021 samples collected from field-edge monitoring wells. Clothianidin concentrations ranged from 0.0125 to 1.63 µg/L and thiamethoxam concentrations ranged from 0.0316 to 3.54 µg/L. These detected concentrations do not exceed any of the respective DHS drinking water health advisories for clothianidin or thiamethoxam.

Imidacloprid concentrations exceeding laboratory reporting limits were detected in 50% of the 2021 groundwater samples collected. It was detected in samples collected from 14 of 24 sites at concentrations ranging from 0.0106 to 2.77 µg/L, an increase in maximum concentration, relative to the maximum of 0.854 µg/L observed in 2020. Imidacloprid exceeded the DHS drinking water health advisory of 0.2 µg/L in 12 samples. These groundwater samples were collected from sites within the Central Sands Agricultural Region and Lower Wisconsin River Valley (Adams, Iowa, Sauk, and Waushara Counties). The imidacloprid data relative to each monitoring location is summarized in [Table B 5 in Appendix B](#).

One observation regarding the 2021 data suggests that the imidacloprid and thiamethoxam are migrating vertically and horizontally within Central Sands Agricultural Region aquifers. Concentrations do not fluctuate seasonally, but greater concentrations have been detected in the groundwater collected from deeper screened wells at sites AD2-5, AD3-3, AD5-5, and WS7-3 compared to adjacent shallow wells. Additionally, imidacloprid and thiamethoxam have also been detected in nearby surface water samples indicating that

groundwater is discharging to surface water year-round as base flow (see DATCP's *2021 Surface Water Sampling Report* - Wisconsin Department of Agriculture, Trade and Consumer Protection, 2023b).

Results from DATCP's Field-Edge Groundwater Monitoring Program can also be compared to nearby historical Surface Water Sampling Program results. This data can then be used to further evaluate mobility, persistence, and discharge to surface water. DATCP intends to report findings of the evaluation along with an evaluation of historical results as part of DATCP's upcoming detailed comprehensive report for each field-edge site.

Alachlor:

As noted previously, alachlor ESA was the third most frequently detected compound in 2021 samples. It was detected in excess of laboratory reporting limits in more than 65% of the samples collected and at 19 of the 24 field-edge monitoring sites. The alachlor ESA data relative to each monitoring location is summarized in [Table B 6](#) in [Appendix B](#).

Alachlor ESA concentrations ranged from 0.0611 to 15.5 µg/L in 2021 samples. The greatest concentration of alachlor ESA was 15.5 µg/L collected from monitoring well JN3-1. This is the first year that an alachlor ESA concentration exceeding the 4.0 µg/L Wis. Admin. Code ch. NR 140 PAL was detected in a field-edge groundwater sample collected at this site.

As observed since 2017, groundwater samples collected from deeper wells AD5-5 and WS7-3 detected alachlor ESA at concentrations in excess of the Wis. Admin. Code ch. NR 140 PAL of 4.0 µg/L. No PAL exceedances were observed in samples collected from wells screened at shallower depths at these same sites between 2018 and 2021. Although alachlor ESA remains at concentrations in excess of the PAL, it cannot be attributed to current use at nearby fields. Alachlor ESA is a breakdown product of alachlor. Alachlor production ceased in December 2014 and could not be sold in Wisconsin after August 2018. The parent alachlor was not detected in excess of laboratory reporting limits in any samples collected in 2021. These results were also observed with samples collected between 2018 and 2020.

Alachlor ESA was also widely detected in surface water and groundwater samples collected throughout the state. Because it is no longer sold in Wisconsin and field use has declined, it is expected that these metabolite concentrations will also decline over time. Additional data collection and evaluation of data from multiple years is needed to validate these observations.

Atrazine:

There are currently 101 atrazine Prohibition Areas (PAs) covering approximately 1.2 million acres within Wisconsin. It is illegal to apply any pesticide containing the active ingredient atrazine within an atrazine PA. In non-PAs, atrazine use is restricted but not prohibited. Since PAs have been in place for 10 years or more, it is anticipated that atrazine and its metabolite concentrations in groundwater would be limited, or not present at all. Of the 24 field-edge sites in the Program, 11 are located within a PA. No grower self-reported atrazine use on adjacent fields within the PAs.

Atrazine or one of its breakdown products (de-ethyl atrazine, de-isopropyl atrazine, and di-amino atrazine) were detected in excess of laboratory reporting limits in 46% of the groundwater samples collected in 2021. No atrazine was detected at concentrations exceeding the 3.0 µg/L Wis. Admin. Code ch. NR 140 ES. However, atrazine TCR was observed in 11 groundwater samples at a concentration greater than the 0.3 µg/L Wisc. Admin. Code ch. NR 140 PAL. Concentrations for atrazine TCR ranged from 0.0507 to 1.297 µg/L. Parent atrazine and metabolite data for each monitoring site is presented in [Table B 7](#) in [Appendix B](#).

The 2021 groundwater results indicated atrazine or one of its metabolites was detected in samples collected from 17 of the 24 sites. Groundwater samples with detections in excess of the Wis. Admin. Code ch. NR 140 PAL for atrazine TCR were collected from monitoring well networks located at eleven of the 24 sites:


- three locations in Adams and Waushara counties;
- two locations in Iowa County; and
- one location in St. Croix County, Sauk County, and Waupaca County.

Of those 11 sites, five are located in a PA: Iowa County (IW1, IW2), St. Croix County (SC1), Sauk County (SK6), and Waushara County (WS4). Of the five locations within a PA, parent material atrazine was found in excess of detection limits at sites IW1, IW2, and SK6. All of these detections were identified in groundwater

samples collected from the new piezometers constructed in 2021 at the deepest monitoring depths. Based on grower self-reporting, atrazine has not been used on the adjacent WS4 fields for over 20 years. These results indicate that the source for the parent material atrazine detections may be older, not from adjacent fields, and beyond the immediate area.

As observed during previous years, the greatest concentrations of atrazine TCR in 2021 samples, were typically detected in samples collected from deeper screened wells. Figure 10 depicts atrazine TCR concentrations relative to groundwater sample well depth. As indicated, elevated concentrations of atrazine TCR were detected in samples collected from monitoring wells screened between 50 and 60 feet below ground surface (bgs), and at deeper wells screened between 80 and 115 feet bgs. On average, shallow wells screened between 10 and 40 feet bgs detected atrazine TCR at lesser concentrations. Based on atrazine TCR concentrations observed across the aquifer depth, it is possible that atrazine is applied at nearby agricultural fields at rates that are not affecting shallow groundwater quality. The greater atrazine concentrations observed at depth likely indicate affects from historic use rather than an on-going source from field use. A trend analysis is needed to show all historical groundwater data to determine if the atrazine TCR concentrations are decreasing within PAs as intended. DATCP intends to report these finding along with an evaluation of historical results as part of DATCP's detailed comprehensive report for each field-edge site.

Figure 10: 2021 Atrazine TCR Concentrations relative to Groundwater Sample Well Depth



Notes: Line through data represents trend of concentrations relative to depth.

Nitrogen:

DATCP's Field-Edge Groundwater Monitoring Program primary focus is on pesticide affects to groundwater quality. In addition to pesticides, BLS includes nitrogen as nitrate plus nitrite analyses. Nitrogen impacts in groundwater and drinking water are the responsibility of WDNR. However, BLS includes nitrogen as nitrate plus nitrite analyses as part of this program, and that data is shared with WDNR.


Nitrogen as nitrate plus nitrite was detected in excess of laboratory reporting limits in 101 of the 106 field-edge groundwater samples collected in 2021. The average nitrogen concentration for all 2021 samples was 16.28 milligram per liter (mg/L or parts per million [ppm]), which is slightly less than last year's (2020) average of concentration of 16.89 ppm. This continues the overall decreasing trend calculated over the past five years as summarized in Table 3.

Table 3: Average Nitrogen as Nitrate plus Nitrite Concentration over Previous Years

| Year | Average Nitrogen-Nitrate/Nitrite Concentration (in parts per million) |
|------|---|
| 2017 | 17.90 |
| 2018 | 17.72 |
| 2019 | 14.61 |
| 2020 | 16.89 |
| 2021 | 16.28 |

The Wis. Admin. Code ch. NR 140 ES of 10 mg/L for nitrogen as nitrate plus nitrite was exceeded in 75 of the 106 groundwater samples collected in 2021. An additional 31 samples exceeded the Wis. Admin. Code ch. NR 140 PAL of 2.0 mg/L. The greatest concentration of nitrogen (41.2 mg/L) was detected in the SK6-2 groundwater sample collected in the fall at a Sauk County station. All nitrogen as nitrate plus nitrite data relative to each monitoring location is summarized in Table B 8 of Appendix B. Figure 11 depicts the 2021 nitrogen concentration distribution.


Figure 11: Nitrogen as Nitrate plus Nitrite Results Distribution in Groundwater Samples from All-Wells



Nitrogen as nitrate plus nitrite concentrations were also compared to wells screened at different depths. Figure 12 depicts nitrogen concentrations for all wells by depth. As indicated, nitrogen as nitrate plus nitrite

was detected over a wide range of concentrations in groundwater samples collected from wells screened at shallow depths (between 10 and 40 feet bgs) compared to deeper wells. Groundwater samples collected from deeper wells typically detected nitrogen as nitrate plus nitrite at greater concentrations. As indicated, nitrogen as nitrate plus nitrite exceeded the 10 mg/L ES in samples collected from nearly all the monitoring wells screened across the aquifer at a depth greater than 40 feet, and in more than half the wells less than 40 feet deep.


Figure 12: 2021 Nitrogen as Nitrate plus Nitrite Concentrations relative to Groundwater Sample Well Depth



Notes: Line through data represents trend of concentrations relative to depth.


Groundwater samples collected from deeper screened wells also show less seasonal variation in nitrogen concentrations compared to shallow wells. As depicted on [Figure 13](#) below, nitrogen as nitrate plus nitrite concentrations fluctuated between -5 mg/L to + 5 mg/L in samples collected between spring and fall 2021 at the majority of monitoring well locations. On average, nitrogen concentrations increased by 0.32 mg/L between spring and fall. Overall, this suggests that nitrogen as nitrate plus nitrite concentrations for the majority of wells indicate little seasonal variation.

Figure 13: 2021 Nitrogen as Nitrate plus Nitrite Concentrations Variability from Spring to Fall at Individual Wells



When seasonal data is plotted based on nitrogen as nitrate - nitrite concentration variances relative to groundwater depths, a relationship does not appear. This likely indicates nitrogen applications at the surface does not influence groundwater quality seasonally. As depicted on the [Figure 14](#) below, groundwater samples collected from shallower wells have a similar range of variability in nitrogen concentrations to deeper wells. Nitrogen as nitrate plus nitrite concentrations in samples collected from deeper screened wells are expected to show less variability and serve as a baseline, where little seasonal influence should be occurring.

Figure 14: 2021 Nitrogen Concentrations Variability by Depth from Spring to Fall of Individual Wells



2022 Program Goals and Objectives

The Field-Edge Groundwater Monitoring Program mission is to monitor groundwater quality at strategic geographic locations within agricultural areas to characterize agrichemical migration to underlying aquifers, and act as an early warning signal for nearby drinking water wells. The Program will continue in 2022.

Program goals for 2022 include:

- Collaborate with BLS and develop a 2022 Field-Edge Groundwater Monitoring Program Sampling Plan.
- Conduct a groundwater sampling event in the spring and fall from the Program’s groundwater monitoring network. This will include continuing to analyze a certain set of samples for glyphosate.
- Document annual activities completed and summarize results for each site in a letter sent to each grower.
- Document the annual activities completed and summarize results in a *2022 Field-Edge Groundwater Monitoring Program Summary Report*.

2022 data will be added to the existing database to ensure that long-term water level and groundwater monitoring data can be used to identify trends in groundwater quality over time. Long-term groundwater quality trends may be used to further evaluate the effectiveness of atrazine Pas. Long-term groundwater data will also be compared to surface water data from within the same watershed to identify potential relationships between surface water and groundwater quality. This evaluation may also be used to evaluate seasonal surface water flow variations and base flow groundwater discharge to surface water. DATCP intends to report finding along with an evaluation of historical results as part of DATCP’s detailed comprehensive report for each field-edge site.

Acknowledgments

ACM's financial information includes the state fiscal year (FY) 2021 and 2022, from July 1, 2020 through June 30, 2022. Federal grants operate October 1 through September 30 (2021 and 2022). This report covers those portions of the federal grants that occurred during the state fiscal year. The primary sources of revenue for ACM are industry fees for licenses, permits, registrations, and tonnage under the feed, fertilizer, soil and plant additive, lime, and pesticide programs. In addition, a federal grant provides some funding to cover annual pesticide program expenses. ACM recognizes these important partnerships with industry and the federal government and works hard to maximize the use of this funding for the benefit of the industry, consumers, and the environment.

The raw data required to reproduce the above findings are available upon request. For any questions and clarifications, please do not hesitate to reach out to us at DATCPGW@wisconsin.gov or at (608) 224-4502.

References

National Centers for Environmental Information (NCEI). (2021). *Annual 2021 National Climate Report*. Retrieved from <https://www.ncei.noaa.gov/access/monitoring/monthly-report/national/202113>

NOAA - National Oceanic and Atmospheric Administration. (2023a). *National Centers for Environmental Information*. Retrieved from Storm Events Database: <https://www.ncdc.noaa.gov/stormevents/>

NOAA - National Oceanic and Atmospheric Administration. (2023b). *National Integrated Drought Information System*. Retrieved from <https://www.drought.gov/>

United States Department of Agriculture - National Agricultural Statistics Service. (2023). *Surveys*. Retrieved from Agricultural Chemical Use Program.

Wisconsin Department of Agriculture, Trade and Consumer Protection. (2023a). *Wisconsin Agricultural Statistics*. Retrieved from <https://datcp.wi.gov/Pages/Publications/WIAgStatistics.aspx>

Wisconsin Department of Agriculture, Trade and Consumer Protection. (2023b). *2021 Surface Water Pesticide Monitoring Program Annual Report*.

Wisconsin Department of Agriculture, Trade and Consumer Protection. (2021). *Sample Collection - Private Well*.

Wisconsin Department of Natural Resources. (1996). *Groundwater Sampling Field Manual*.

Wisconsin State Climatology Office. (2023). Retrieved from Wisconsin Climate Watch Archive: <https://www.aos.wisc.edu/~sco/clim-watch/archives.html#monthly2021>

Appendix A

The acronyms and terminology included on this list are generic definitions intended to help understand the Field-Edge Monitoring Program. Some of these terms are more specifically defined in various regulations.

ACRONYMS

| | |
|------------------------|--|
| µg/l _____ | Micrograms per liter (a liquid equivalent of ppb) |
| ACM _____ | DATCP Bureau of Agrichemical Management |
| AMPA _____ | Aminomethylphosphonic acid |
| Bgs _____ | Below ground surface |
| BLS _____ | DATCP Bureau of Laboratory Services |
| CAS _____ | Chemical Abstract Service |
| CIT _____ | clothianidin, imidacloprid and thiamethoxam |
| DATCP _____ | Department of Agriculture, Trade and Consumer Protection |
| DADK _____ | Desaminodiketo |
| ES _____ | Enforcement Standard |
| ESA _____ | Ethane Sulfonic Acid |
| GC _____ | Gas Chromatography |
| GCC _____ | Wisconsin Groundwater Coordinating Council |
| HARS _____ | Hancock Agricultural Research Station |
| ISO _____ | International Organization for Standardization |
| LC _____ | Liquid Chromatography |
| mg/L _____ | Milligrams per liter (a liquid equivalent of ppm) |
| MS _____ | Mass Spectroscopy |
| msl _____ | Mean sea level |
| N _____ | Nitrogen |
| ND _____ | No Detect - concentrations are less than laboratory reporting limits |
| NOAA _____ | National Oceanic and Atmospheric Administration |
| OA _____ | Oxanilic Acid |
| PA _____ | Prohibition Area |
| PAL _____ | Preventive Action Limit |
| PPB _____ | Parts per billion |
| PPM _____ | Parts per million |
| TCR _____ | Total chlorinated residues of atrazine |
| TPVC _____ | Top of well casing |
| TSAMP _____ | Targeted Sampling Program |
| USDA _____ | U.S. Department of Agriculture |
| WDHS _____ | Wisconsin Department of Health Services |
| WDNR _____ | Wisconsin Department of Natural Resources |
| WGNHS _____ | Wisconsin Geological and Natural History Survey |
| Wis. Admin. Code _____ | Wisconsin Administrative Code |
| WUWN _____ | Wisconsin Unique Well Number |
| US EPA _____ | United States - Environmental Protection Agency |
| USDA _____ | United States Department of Agriculture |

DEFINITIONS

Analyte - A chemical substance that has a defined Chemical Abstract Service (CAS) number

Atrazine Prohibition Area - An area where atrazine use is currently prohibited under Administrative Code ATCP 30

Chronic Exposure value - The highest concentration of a chemical to which the organism can be exposed without causing chronic toxicity to the organism in question

Compound - A substance formed by the chemical union of two or more ingredients

Detection - When an analyte has a concentration that can be quantified (i.e., a concentration greater than the Laboratory Reporting Limit)

Herbicide - A pesticide used to kill or inhibit the growth of plants, weeds, or grasses

Insecticide - A pesticide used to kill or inhibit the growth of insects

Metabolite or Residual compound or Breakdown product - A chemical substance left behind by a parent compound that has degraded through natural chemical breakdown and/or been metabolized by bacteria

Neonicotinoids - Insecticides that target the neurological systems of insects. The neonicotinoid family includes acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, nithiazine, thiacloprid, and thiamethoxam

NR140 - Wisconsin Administrative Code which establishes groundwater quality standards and required responses when the standards are exceeded

Pesticide - Substance used to kill, repel, or control certain forms of plant or animal life that are considered to be pests. The pesticide category includes herbicides, insecticides, rodenticides, fungicides, and bactericides

Reporting limit - The minimum analyte concentration that can be reliably quantified and reported by the laboratory

Total chlorinated residues (TCR) of atrazine - Sum of atrazine and atrazine metabolites (de-ethyl atrazine, de-isopropyl atrazine, and diamino atrazine)

APPENDIX B

Table B 1: Field-Edge Groundwater Monitoring Program - Monitoring Wells and Piezometers Construction Specifications

| County | Site (Grower) | Well Identification | WUWN | Year Constructed | Prohibition Area | Irrigation Available | Ground Elevation (MSL) | TPVC Elevation (MSL) | Well Depth (ft) | Bottom of Well (MSL) | Screen Length (ft) | Top of Screen (ft) | Sampling Method | | | |
|-----------|---------------|---------------------|-------|------------------|------------------|----------------------|------------------------|----------------------|-----------------|----------------------|--------------------|--------------------|---------------------------------|-----------------|--------|-----------------|
| Adams | AD2 | AD2-1 | BH954 | 1987 | No | Yes | 1,051.7 | 1,053.96 | 17.87 | 1,036.09 | 5 | 1,053.96 | Peristolic Pump | | | |
| | | AD2-2 | BH953 | 1987 | | | | 1,054.14 | 22.83 | 1,031.31 | 5 | 1,054.14 | | | | |
| | | AD2-3 | BH952 | 1987 | | | | 1,054.17 | 27.62 | 1,026.55 | 5 | 1,054.17 | | | | |
| | | AD2-4 | VR844 | 2017 | | | | 1,054.44 | 54.70 | 999.74 | 5 | 1,054.44 | Whale Pump and Dedicated Tubing | | | |
| | | AD2-5 | VR845 | 2017 | | | | 1,054.35 | 85.70 | 968.65 | 5 | 1,054.35 | | | | |
| | | AD2-6 | PT421 | 2021 | | | | -- | 116.40 | -- | 5 | -- | | | | |
| | AD3 | AD3-1 | BH999 | 1987 | No | Yes | 1,008.0 | 1,010.48 | 14.93 | 995.55 | 5 | 1,010.48 | | | | |
| | | AD3-2 | B000 | 1987 | | | | 1,010.34 | 19.64 | 990.70 | 5 | 1,010.34 | | | | |
| | | AD3-3 | B001 | 1987 | | | | 1,010.44 | 24.69 | 985.75 | 5 | 1,010.44 | | | | |
| | AD4 | AD4-1 | BH996 | 1987 | No | Yes | 1,013.9 | 1,017.38 | 24.71 | 992.67 | 5 | 1,017.38 | | | | |
| | | AD4-2 | BH997 | 1987 | | | | 1,017.26 | 29.69 | 987.57 | 5 | 1,017.26 | | | | |
| | | AD4-3 | BH998 | 1987 | | | | 1,016.56 | 34.57 | 981.99 | 5 | 1,016.56 | | | | |
| | AD5 | AD5-1 | AD5-1 | CL461 | 1988 | No | Yes | 1,051.1 | 1,053.18 | 15.24 | 1,037.94 | 5 | 1,053.18 | Peristolic Pump | | |
| | | | AD5-2 | CL455 | 1988 | | | | 1,053.31 | 19.91 | 1,033.40 | 5 | 1,053.31 | | | |
| | | | AD5-3 | CL456 | 1988 | | | | 1,053.27 | 25.23 | 1,028.04 | 5 | 1,053.27 | | | |
| AD5-4 | | AD5-4 | VR846 | 2017 | 1,053.63 | | | | 53.20 | 1,000.43 | 5 | 1,053.63 | Whale Pump and Dedicated Tubing | | | |
| | | AD5-5 | VR847 | 2017 | 1,053.68 | | | | 85.70 | 967.98 | 5 | 1,053.68 | | | | |
| | | AD5-6 | PT422 | 2021 | -- | | | | 117.50 | -- | 5 | -- | | | | |
| Barron | BR3 | BR3-1 | BR279 | 1987 | No | Yes | 1,052.7 | 1,055.79 | 15.03 | 1,040.76 | 5 | 1,055.79 | Peristolic Pump | | | |
| | | BR3-2 | BR280 | 1987 | | | | 1,055.37 | 20.02 | 1,035.35 | 5 | 1,055.37 | | | | |
| | | BR3-3 | BR281 | 1987 | | | | 1,054.93 | 25.02 | 1,029.91 | 5 | 1,054.93 | | | | |
| | | BR3-4 | BR250 | 1985 | | | | 744.38 | 12.10 | 732.28 | 5 | 744.38 | | | | |
| Dane | DN1 | DN1-1 | PT428 | 2021 | 93-57-04 | Yes | 743.7 | 745.32 | 14.90 | 730.42 | 5 | 745.32 | Dedicated Bailer | | | |
| | | DN1-2 | BR251 | 1985 | | | | 745.87 | 17.40 | 728.47 | 5 | 745.87 | Peristolic Pump | | | |
| | | DN1-3 | BR252 | 1985 | | | | 746.08 | 22.20 | 723.88 | 5 | 746.08 | | | | |
| | | DN1-4 | BR253 | 1985 | | | | 746.08 | 22.20 | 723.88 | 5 | 746.08 | | | | |
| Dunn | DU1 | DU1-1 | AO384 | 1989 | No | Yes | 852.5 | 853.92 | 34.90 | 819.02 | 5 | 853.92 | Dedicated Bailer | | | |
| | | DU1-2 | AO385 | 1989 | | | | 854.87 | 40.80 | 814.07 | 5 | 854.87 | | | | |
| | | DU1-3 | AO386 | 1989 | | | | 855.12 | 46.10 | 809.02 | 5 | 855.12 | | | | |
| | DU2 | DU2-1 | AO387 | 1989 | | | | No | Yes | 856.2 | 858.05 | 26.70 | 831.35 | 5 | 858.05 | Peristolic Pump |
| | | DU2-2 | AO388 | 1989 | | | | | | | 858.17 | 31.30 | 826.87 | 5 | 858.17 | |
| | | DU2-3 | AO389 | 1989 | | | | | | | 858.48 | 36.60 | 821.88 | 5 | 858.48 | |
| Grant | GR1 | GR1-1 | BR255 | 1985 | 93-57-04 | No | 683.8 | 686.32 | 12.50 | 673.82 | 5 | 686.32 | Peristolic Pump | | | |
| | | GR1-2 | BR256 | 1985 | | | | 686.48 | 17.30 | 669.18 | 5 | 686.48 | | | | |
| | | GR1-3 | BR257 | 1985 | | | | 686.12 | 21.60 | 664.52 | 5 | 686.12 | | | | |
| | | GR1-4 | BR258 | 1985 | | | | 686.12 | 21.60 | 664.52 | 5 | 686.12 | | | | |
| Iowa | IW1 | IW1-1 | BH955 | 1986 | 93-57-04 | Yes | 722.5 | 723.85 | 17.10 | 706.75 | 5 | 723.85 | Peristolic Pump | | | |
| | | IW1-2 | BH956 | 1986 | | | | 723.84 | 21.30 | 702.54 | 5 | 723.84 | | | | |
| | | IW1-3 | BH957 | 1986 | | | | 723.67 | 26.70 | 696.97 | 5 | 723.67 | | | | |
| | | IW1-4 | BR259 | 1986 | | | | 723.67 | 61.99 | 661.68 | 5 | 723.67 | Whale Pump and Dedicated Tubing | | | |
| | | IW1-5 | BR260 | 1986 | | | | 723.06 | 93.97 | 629.09 | 5 | 723.06 | | | | |
| | | IW1-6 | BR261 | 1986 | | | | 726.76 | 14.80 | 711.96 | 5 | 726.76 | Peristolic Pump | | | |
| | | IW1-7 | BH967 | 1987 | | | | 726.50 | 19.70 | 706.80 | 5 | 726.50 | | | | |
| | | IW1-8 | PT425 | 2021 | | | | 726.40 | 24.70 | 701.70 | 5 | 726.40 | | | | |
| | IW2 | IW2-1 | BR036 | 1986 | 93-57-04 | Yes | 723.8 | 725.89 | 65.92 | 659.97 | 5 | 725.89 | Whale Pump and Dedicated Tubing | | | |
| | | IW2-2 | BR037 | 1986 | | | | 726.24 | 94.81 | 631.43 | 5 | 726.24 | | | | |
| Jackson | JK3 | JK3-1 | JH991 | 2005 | 94-27-04 | No | 1,025.3 | 1,028.06 | 27.33 | 1,000.73 | 10 | 1,028.06 | Peristolic Pump | | | |
| | | JK3-2 | JH981 | 2005 | | | | 1,023.7 | 1,026.44 | 25.77 | 1,000.67 | 10 | | 1,026.44 | | |
| Juneau | JN1 | JN1-1 | BR046 | 1985 | No | Yes | 939.7 | 941.26 | 11.70 | 929.56 | 5 | 941.26 | Peristolic Pump | | | |
| | | JN1-2 | BR047 | 1985 | | | | 941.21 | 16.70 | 924.51 | 5 | 941.21 | | | | |
| | | JN1-3 | BR048 | 1985 | | | | 941.34 | 21.50 | 919.84 | 5 | 941.34 | | | | |
| | JN3 | JN3-1 | JH937 | 2005 | | | | 94-29-01 | No | 901.5 | 903.84 | 20.42 | 883.42 | 10 | 903.84 | Peristolic Pump |
| JN3-2 | | JH936 | 2005 | 902.0 | 905.20 | 18.14 | 887.06 | | | | 10 | 905.20 | | | | |
| La Crosse | LC2 | LC2-1 | VZ391 | 2011 | No | Yes | 684.1 | 686.40 | 49.22 | 637.18 | 10 | 686.40 | Dedicated Bailer | | | |
| | | LC2-2 | VZ392 | 2011 | | | | 687.8 | 681.91 | 43.98 | 637.93 | 10 | | 681.91 | | |

| | | | | | | | | | | | | | |
|-------------|-----|-----------|-------|------|----------|-----|----------|----------|--------|----------|----------|---------------------------------|---------------------------------|
| Langlade | LN1 | LN1-1 | BH964 | 1986 | No | No | 1,471.6 | 1,473.85 | 14.80 | 1,459.05 | 5 | 1,473.85 | Peristolic Pump |
| | | LN1-2 | BH965 | 1986 | | | | 1,474.44 | 19.70 | 1,454.74 | 5 | 1,474.44 | |
| | | LN1-3 | BH966 | 1986 | | | | 1,473.74 | 24.80 | 1,448.94 | 5 | 1,473.74 | |
| Portage | PR1 | PR1-1 | BR207 | 1986 | No | Yes | 1,079.7 | 1,082.01 | 12.70 | 1,069.31 | 5 | 1,082.01 | Peristolic Pump |
| | | PR1-2 | BR208 | 1988 | | | | 1,081.94 | 17.60 | 1,064.34 | 5 | 1,081.94 | |
| | | PR1-3 | BR209 | 1988 | | | | 1,081.72 | 22.50 | 1,059.22 | 5 | 1,081.72 | Whale Pump and Dedicated Tubing |
| | | PR1-4 | VR848 | 2017 | | | | 1,082.83 | 55.30 | 1,027.53 | 5 | 1,082.83 | |
| | | PR1-5 | VR849 | 2017 | | | | 1,082.77 | 84.70 | 998.07 | 5 | 1,082.77 | |
| St. Croix | SC1 | SC1-1 | JH938 | 2005 | 94-56-02 | Yes | 1,006.8 | 1,010.14 | 24.87 | 985.27 | 10 | 1,010.14 | Peristolic Pump |
| | | SC1-1 (D) | VZ390 | 2011 | | | | 1,009.16 | 30.10 | 979.06 | 10 | 1,009.16 | |
| | | SC1-2 | JH939 | 2005 | | | 1,003.9 | 21.87 | 984.76 | 10 | 1,006.63 | | |
| | | SC1-2(D) | VZ393 | 2011 | | | 1,006.40 | 30.17 | 976.23 | 10 | 1,006.40 | | |
| Sauk | SK6 | SK6-1 | BB246 | 1988 | 93-57-04 | Yes | 711.8 | 713.68 | 14.92 | 698.76 | 5 | 713.68 | Peristolic Pump |
| | | SK6-2 | BB247 | 1988 | | | | 713.37 | 20.04 | 693.33 | 5 | 713.37 | |
| | | SK6-3 | BB248 | 1988 | | | 713.55 | 25.10 | 688.45 | 5 | 713.55 | Whale Pump and Dedicated Tubing | |
| | | SK6-4 | PT424 | 2021 | | | 710.2 | 711.56 | 53.42 | 658.14 | 5 | | 711.56 |
| Trempealeau | TR1 | TR1-1 | PX201 | 2005 | No | Yes | 730.4 | 733.29 | 75.55 | 657.74 | 10 | 733.29 | Dedicated Bailer |
| | | TR1-2 | PX202 | 2005 | | | 731.1 | 733.83 | 75.20 | 658.63 | 10 | 733.83 | |
| Waupaca | WP2 | WP2-1 | JH985 | 2005 | 94-69-01 | No | 908.4 | 911.03 | 20.45 | 890.58 | 10 | 911.03 | Peristolic Pump |
| | | WP2-2 | JH984 | 2005 | | | 905.7 | 908.82 | 20.43 | 888.39 | 10 | 908.82 | |
| Waushara | WS4 | WS4-1 | BB258 | 1988 | 93-70-01 | Yes | 1,082.4 | 1,084.97 | 17.13 | 1,067.84 | 5 | 1,084.97 | Peristolic Pump |
| | | WS4-2 | BB259 | 1988 | | | | 1,085.03 | 22.02 | 1,063.01 | 5 | 1,085.03 | |
| | | WS4-3 | BB260 | 1988 | | | | 1,084.98 | 27.16 | 1,057.82 | 5 | 1,084.98 | |
| | | WS4-4 | BB261 | 1988 | | | | 1,084.88 | 31.94 | 1,052.94 | 5 | 1,084.88 | |
| | WS6 | WS6-1 | JH989 | 2005 | 93-70-01 | Yes | 1,076.8 | 1,080.90 | 18.27 | 1,062.63 | 10 | 1,080.90 | Peristolic Pump |
| | | WS6-2 | JH990 | 2005 | | | | 1,079.07 | 17.02 | 1,062.05 | 10 | 1,079.07 | |
| | WS7 | WS7-1 | VR841 | 2017 | No | Yes | 1,075.7 | 1,078.65 | 18.40 | 1,060.25 | 10 | 1,078.65 | Peristolic Pump |
| | | WS7-2 | VR842 | 2017 | | | | 1,078.79 | 54.70 | 1,024.09 | 5 | 1,078.79 | |
| | | WS7-3 | VR843 | 2017 | | | | 1,078.78 | 84.80 | 993.98 | 5 | 1,078.78 | Whale Pump and Dedicated Tubing |
| | | WS7-4 | PT423 | 2021 | | | | -- | 104.10 | -- | 5 | -- | |







- Notes:
- Elevation surveying in progress.
 - 1 Monitoring well was abandoned on May 30, 2019 because integrity of protective casing was compromised during spring 2019 sampling.
 - 2 Monitoring well was abandoned on December 13, 2018 because integrity of protective casing was compromised by a vehicle prior to fall 2018 sampling.
 - 3 Monitoring wells were abandoned June 11, 1993 because they were no longer needed for the monitoring program.
 - 4 Monitoring wells were abandoned December 1, 2021 because ownership no longer wished to participate in the monitoring program.
 - WUWN Wisconsin Unique Well Number
 - MSL Mean sea level
 - TPVC Top of well casing (PVC)
 -  Monitoring Well/Piezometer abandoned.
 -  Monitoring Well/Piezometer construction was financed by a 2021 U.S. EPA grant.
 -  Monitoring Well/Piezometer construction was financed by a 2017 U.S. EPA grant.
 -  Monitoring Well/Piezometer construction was financed by a 2011 U.S. EPA grant.
 -  Monitoring Well/Piezometer construction was financed by a 2005 U.S. EPA grant.
 -  Monitoring Wells/Piezometers associated with initial program activities and financed by State.

Table B 2: 2021 Sample Analytes, Applicable Wis. Admin. Code ch. NR 140 PALs & ESs, Drinking Water Health Advisories, and Reporting Limits

| Analyte Description | PAL (µg/L) | ES (µg/L) | Advisory* | Reporting Limit (µg/l) |
|--|------------------|------------------|-----------------|------------------------|
| 2,4-D (dichlorophenoxyacetic acid) | 7 | 70 | | 0.050 |
| 2,4-DB | | | | 0.80 |
| 2,4-DP | | | | 0.050 |
| 2,4,5-T | | | | 0.050 |
| 2,4,5-TP (trichlorophenoxy-prop. acid) | 5 | 50 | | 0.050 |
| ACETAMIPRID | | | | 0.010 |
| ACETOCHLOR | 0.7 | 7 | | 0.050 |
| ACETOCHLOR ESA | 46 ¹ | 230 ¹ | | 0.050 |
| ACETOCHLOR OA | 46 ¹ | 230 ¹ | | 0.30 |
| ACIFLUORFEN | | | | 0.050 |
| ALACHLOR | 0.2 | 2 | | 0.050 |
| ALACHLOR ESA | 4 | 20 | | 0.053 |
| ALACHLOR OA | | | | 0.25 |
| ALDICARB SULFONE | | | | 0.050 |
| ALDICARB SULFOXIDE | | | | 0.071 |
| AMINOPYRALID | | | | 0.150 |
| ATRAZINE | 0.3 | 3 | | 0.050 |
| DE-ETHYL ATRAZINE | 0.3 | 3 | | 0.050 |
| DEISOPROPYL ATRAZINE | 0.3 | 3 | | 0.050 |
| DIAMINO ATRAZINE | 0.3 | 3 | | 0.20 |
| ATRAZINE TCR (calculated) | 0.3 ³ | 3 ³ | | 0.050 |
| AZOXYSTROBIN | | | | 0.050 |
| BENFLURALIN | | | | 0.050 |
| BENTAZON | 60 | 300 | | 0.050 |
| BICYCLOPYRONE | | | | 0.050 |
| BROMACIL | | | | 0.050 |
| BIFENTHRIN | | | | 0.005 |
| CARBARYL | 4 | 40 | | 0.050 |
| CARBOFURAN | 8 | 40 | | 0.050 |
| CHLORAMBEN | 30 | 150 | | 0.32 |
| CHLORANTRANILIPROLE | | | 16,000 | 0.050 |
| CHLOROTHALONIL | | | | 0.10 |
| CHLORPYRIFOS | 0.4 | 2 | | 0.050 |
| CHLORPYRIFOS OXYGEN ANALOG | | | | 0.050 |
| CLOMAZONE | | | | 0.050 |
| CLOPYRALID | | | | 0.050 |
| CLOTHIANIDIN | | | 1,000 | 0.010 |
| CYANTRANILIPROLE | | | | 0.050 |
| CYCLANILIPROLE | | | | 0.20 |
| CYFLUTHRIN | | | | 0.050 |
| CYPERMETHRIN | | | | 0.10 |
| CYPROSULFAMIDE | | | | 0.050 |
| DACTHAL | 14 | 70 | | 0.050 |
| DACTHAL DI-ACID | | | 70 ² | 0.050 |
| DACTHAL MONO-ACID | | | 70 ² | 0.050 |
| DIAZINON | | | | 0.050 |
| DIAZINON OXYGEN ANALOG | | | | 0.050 |
| DICAMBA | 60 | 300 | | 0.30 |
| DICHOBENIL | | | | 0.050 |
| DIMETHENAMID | 5 | 50 | | 0.050 |
| DIMETHENAMID ESA | | | | 0.050 |
| DIMETHENAMID OA | | | | 0.050 |
| DIMETHOATE | 0.4 | 2 | | 0.050 |
| DINOTEFURAN | | | | 0.010 |
| DIURON | | | | 0.050 |

| Analyte Description | PAL (µg/L) | ES (µg/L) | Advisory* | Reporting Limit (µg/l) |
|---------------------------------|------------------|-------------------|----------------|------------------------|
| EPTC | 50 | 250 | | 0.050 |
| ESFENVALERATE | | | | 0.025 |
| ETHALFLURALIN | | | | 0.050 |
| ETHOFUMESATE | | | | 0.050 |
| FLUMETSULAM | | | 10,000 | 0.050 |
| FLUPYRADIFURONE | | | | 0.050 |
| FLUROXYPYR | | | | 0.070 |
| FOMESAFEN | | | 25 | 0.050 |
| GLYPHOSATE | | | 10,000 | 0.50 |
| GLYPHOSATE AMMONIUM | | | | 0.50 |
| AMPA | | | 10,000 | 0.50 |
| HALOSULFURON METHYL | | | | 0.050 |
| HEXAZINONE | | | 400 | 0.050 |
| IMAZAPYR | | | | 0.050 |
| IMAZETHAPYR | | | | 0.050 |
| IMIDACLOPRID | | | 0.2 | 0.010 |
| ISOXAFLOTOLE | | | 3 ⁴ | 0.050 |
| ISOXAFLOTOLE DKN | | | 3 ⁴ | 0.050 |
| LAMBDA-CYHALOTHRIN | | | | 0.020 |
| LINURON | | | | 0.050 |
| MALATHION | | | | 0.050 |
| MCPA | | | | 0.050 |
| MCPB | | | | 0.10 |
| MCPP | | | | 0.050 |
| MESOTRIONE | | | | 0.10 |
| METALAXYL | | | 800 | 0.050 |
| METHYL PARATHION | | | | 0.050 |
| METOLACHLOR | 10 | 100 | | 0.050 |
| METOLACHLOR ESA | 260 ⁵ | 1300 ⁵ | | 0.050 |
| METOLACHLOR OA | 260 ⁵ | 1300 ⁵ | | 0.27 |
| METRIBUZIN | 14 | 70 | | 0.050 |
| METRIBUZIN DA | | | | 0.10 |
| METRIBUZIN DADK | | | | 0.12 |
| METSULFURON-METHYL | | | | 0.050 |
| NICOSULFURON | | | | 0.050 |
| NORFLURAZON | | | | 0.050 |
| OXADIAZON | | | | 0.050 |
| PENDIMETHALIN | | | | 0.050 |
| PERMETHRIN | | | | 0.030 |
| PICLORAM | 100 | 500 | | 0.050 |
| PROMETONE | 20 | 100 | | 0.050 |
| PROMETRYN | | | | 0.050 |
| PROPICONAZOLE | | | | 0.050 |
| PROTHIOCONAZOLE-DESTHIO | | | | 0.050 |
| SAFLUFENACIL | | | 460 | 0.050 |
| SIMAZINE | 0.4 | 4 | | 0.050 |
| SULFENTRAZONE | | | 1,000 | 0.050 |
| SULFOMETURON-METHYL | | | | 0.050 |
| TEBUPIRIMPHOS | | | | 0.050 |
| TEMBOTRIONE | | | | 0.10 |
| THIACLOPRID | | | | 0.010 |
| THIAMETHOXAM | | | 120 | 0.010 |
| THIENCARBAZONE-METHYL | | | 800 | 0.050 |
| TRICLOPYR | | | | 0.050 |
| TRIFLURALIN | 0.75 | 7.5 | | 0.050 |
| NITROGEN-NITRATE/NITRITE (mg/L) | 2 | 10 | | 0.5 |

* Wisconsin Department of Health Services Drinking Water Health Advisory (June 2019, November 2020, Revised February 2022).

¹ Combined sum of acetochlor metabolites ESA and OA.

² Combined sum of metabolites (di- and mono-acid) and parent material dacthal.

³ Total Chlorinated Residue for Atrazine. Combined sum of metabolites (de-ethyl, de-isopropyl) and di-amino) and parent material atrazine.

⁴ Combined sum of metabolite (DKN) and parent material isoxaflutole.

⁵ Combined sum of metolachlor metabolites ESA and OA.

µg/L - micrograms per liter or parts per billion.

mg/L - milligrams per liter or parts per million.

DKN - diketonitrile

ESA - ethane sulfonic acid.

OA - oxanilic acid, can also be identified as OXA.

Table B 3: Field-Edge Groundwater Monitoring Program - 2021 Groundwater Analytical Results

| 2021 Ground Water Project Results (all concentrations in ug/L) | | | | | | | Wisconsin Department of Health Services | Wisconsin Admin. Code Chapter NR 140 | |
|--|-----------------|-----------------|---|--------------------------------------|---------------------------------|---------------------|---|--------------------------------------|-------------------------|
| Pesticide Name | Pesticide Class | Reporting Limit | Number of Sites with Detects ¹ | Number of Total Detects ² | Percent of Samples with Detects | Concentration Range | Drinking Water Health Advisory ³ | Enforcement Standard | Preventive Action Limit |
| 2,4-D (dichlorophenoxyacetic acid) | Herbicide | 0.05 | -- | -- | -- | -- | -- | 70 | 7 |
| 2,4-DB | Herbicide | 1.50 | -- | -- | -- | -- | -- | -- | -- |
| 2,4-DP | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| 2,4,5-T | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| 2,4,5-TP (trichlorophenoxy-propionic acid) | Herbicide | 0.05 | -- | -- | -- | -- | -- | 50 | 5 |
| Acetamiprid | Insecticide | 0.010 | -- | -- | -- | -- | -- | -- | -- |
| Acetochlor | Herbicide | 0.05 | -- | -- | -- | -- | -- | 7 | 0.7 |
| Acetochlor ESA | Metabolite | 0.05 | 11 | 39 | 36.8% | 0.0581-2.32 | -- | 230 ⁴ | 46 ⁴ |
| Acetochlor OA | Metabolite | 0.3 | 1 | 1 | 0.9% | 0.492 | -- | 230 ⁴ | 46 ⁴ |
| Acifluorfen | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Alachlor | Herbicide | 0.05 | -- | -- | -- | -- | -- | 2 | 0.2 |
| Alachlor ESA | Metabolite | 0.053 | 19 | 69 | 65.1% | 0.0611-15.5 | -- | 20 | 4 |
| Alachlor OA | Metabolite | 0.25 | 5 | 7 | 6.6% | 0.251-4.07 | -- | -- | -- |
| Aldicarb Sulfone | Insecticide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Aldicarb Sulfoxide | Insecticide | 0.071 | -- | -- | -- | -- | -- | -- | -- |
| Aminopyralid | Herbicide | 0.15 | -- | -- | -- | -- | -- | -- | -- |
| Atrazine | Herbicide | 0.05 | 8 | 19 | 17.9% | 0.0536-0.328 | -- | 3 | 0.3 |
| De-ethyl atrazine | Metabolite | 0.05 | 13 | 36 | 34.0% | 0.0529-0.744 | -- | 3 | 0.3 |
| De-isopropyl atrazine | Metabolite | 0.05 | 11 | 25 | 23.6% | 0.0507-0.605 | -- | 3 | 0.3 |
| Di-amino atrazine | Metabolite | 0.2 | 10 | 13 | 12.3% | 0.201-0.512 | -- | 3 | 0.3 |
| Atrazine (TCR) | Sumation | 0.05 | 17 | 49 | 46.2% | 0.0507-1.297 | -- | 3 | 0.3 |
| Azoxystrobin | Fungicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Benfluralin | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Bentazon | Herbicide | 0.05 | 6 | 15 | 14.2% | 0.0694-30.5 | -- | 300 | 60 |
| Bicyclopyrone | Herbicide | 0.05 | 2 | 2 | 1.9% | 0.0736-0.0802 | -- | -- | -- |
| Bifentrin | Insecticide | 0.0050 | -- | -- | -- | -- | -- | -- | -- |
| Bromacil | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Carbaryl | Insecticide | 0.05 | -- | -- | -- | -- | -- | 40 | 4 |
| Carbofuran | Insecticide | 0.05 | -- | -- | -- | -- | -- | 40 | 8 |

| | | | | | | | | | |
|---------------------|-------------|-------|----|----|-------|--------------|-----------------|-----|-----|
| Chloramben | Herbicide | 0.32 | -- | -- | -- | -- | -- | 150 | 30 |
| Chlorantraniliprole | Insecticide | 0.050 | 9 | 30 | 28.3% | 0.0696-1.08 | 16,000 | -- | -- |
| Chlorothalonil | Fungicide | 0.10 | -- | -- | -- | -- | -- | -- | -- |
| Chlorpyrifos | Insecticide | 0.05 | -- | -- | -- | -- | -- | 2 | 0.4 |
| Chlorpyrifos Oxon | Metabolite | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Clomazone | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Clopyralid | Herbicide | 0.05 | 1 | 1 | 0.9% | 0.32 | -- | -- | -- |
| Clothianidin | Insecticide | 0.010 | 20 | 79 | 74.5% | 0.0125-1.63 | 1,000 | -- | -- |
| Cyantraniliprole | Insecticide | 0.050 | 4 | 5 | 4.7% | 0.0515-0.224 | -- | -- | -- |
| Cyclaniliprole | Insecticide | 0.2 | -- | -- | -- | -- | -- | -- | -- |
| Cyfluthrin | Insecticide | 0.050 | -- | -- | -- | -- | -- | -- | -- |
| lambda- Cyhalothrin | Insecticide | 0.020 | -- | -- | -- | -- | -- | -- | -- |
| Cypermethrin | Insecticide | 0.1 | -- | -- | -- | -- | -- | -- | -- |
| Cyprosulfamide | Safener | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Dacthal | Herbicide | 0.05 | -- | -- | -- | -- | -- | 70 | 14 |
| Dacthal Di-acid | Metabolite | 0.05 | -- | -- | -- | -- | 70 ⁵ | -- | -- |
| Dacthal Mono-acid | Metabolite | 0.05 | -- | -- | -- | -- | 70 ⁵ | -- | -- |
| Diazinon | Insecticide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Diazinon oxon | Metabolite | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Dicamba | Herbicide | 0.60 | -- | -- | -- | -- | -- | 300 | 60 |
| Dichlobenil | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Dimethenamid | Herbicide | 0.05 | 1 | 1 | 0.9% | 0.214 | -- | 50 | 5 |
| Dimethenamid ESA | Metabolite | 0.05 | 5 | 12 | 11.3% | 0.0685-6.36 | -- | -- | -- |
| Dimethenamid OA | Metabolite | 0.05 | 2 | 2 | 1.9% | 0.086-1.08 | -- | -- | -- |
| Dimethoate | Insecticide | 0.050 | -- | -- | -- | -- | -- | 2 | 0.4 |
| Dinotefuran | Insecticide | 0.010 | -- | -- | -- | -- | -- | -- | -- |
| Diuron | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| EPTC | Herbicide | 0.05 | -- | -- | -- | -- | -- | 250 | 50 |
| Esfenvalerate | Insecticide | 0.025 | -- | -- | -- | -- | -- | -- | -- |
| Ethalfuralin | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Ethofumesate | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Flumetsulam | Herbicide | 0.05 | 2 | 3 | 2.8% | 0.0506-0.159 | 10,000 | -- | -- |
| Flupyradifurone | Insecticide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Fluroxypyr | Insecticide | 0.070 | -- | -- | -- | -- | -- | -- | -- |
| Fomesafen | Herbicide | 0.05 | 3 | 8 | 7.5% | 0.0506-1.23 | 25 | -- | -- |

| | | | | | | | | | |
|-------------------------|-------------|-------|----|-----|-------|-------------|----------------|--------------------|------------------|
| Glyphosate | Herbicide | 0.5 | -- | -- | -- | -- | 10,000 | -- | -- |
| Glyphosate Ammonium | Metabolite | 0.5 | -- | -- | -- | -- | -- | -- | -- |
| AMPA | Metabolite | 0.5 | -- | -- | -- | -- | 10,000 | -- | -- |
| Halosulfuron methyl | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Hexazinone | Herbicide | 0.05 | -- | -- | -- | -- | 400 | -- | -- |
| Imazapyr | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Imazethapyr | Herbicide | 0.05 | 1 | 2 | 1.9% | 0.117-0.228 | -- | -- | -- |
| Imidacloprid | Insecticide | 0.010 | 14 | 53 | 50.0% | 0.0106-2.77 | 0.2 | -- | -- |
| Isoxaflutole | Herbicide | 0.05 | -- | -- | -- | -- | 3 ⁶ | -- | -- |
| Isoxaflutole DKN | Metabolite | 0.05 | -- | -- | -- | -- | 3 ⁶ | -- | -- |
| Linuron | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| MCPA | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| MCPB | Herbicide | 0.1 | -- | -- | -- | -- | -- | -- | -- |
| MCPP | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Malathion | Insecticide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Mesotrione | Herbicide | 0.1 | -- | -- | -- | -- | -- | -- | -- |
| Metalaxyl | Fungicide | 0.05 | 13 | 35 | 33.0% | 0.05-1.43 | 800 | -- | -- |
| Methyl Parathion | Insecticide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Metolachlor | Herbicide | 0.05 | 15 | 45 | 42.5% | 0.0539-5.08 | -- | 100 | 10 |
| Metolachlor ESA | Metabolite | 0.05 | 24 | 104 | 98.1% | 0.0883-34.9 | -- | 1,300 ⁷ | 260 ⁷ |
| Metolachlor OA | Metabolite | 0.27 | 17 | 67 | 63.2% | 0.313-27 | -- | 1,300 ⁷ | 260 ⁷ |
| Metribuzin | Herbicide | 0.05 | 11 | 41 | 38.7% | 0.0738-8.73 | -- | 70 | 14 |
| Metribuzin DA | Metabolite | 0.1 | 6 | 13 | 12.3% | 0.106-0.911 | -- | -- | -- |
| Metribuzin DADK | Metabolite | 0.12 | 11 | 35 | 33.0% | 0.12-5.88 | -- | -- | -- |
| Metsulfuron methyl | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Nicosulfuron | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Norflurazon | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Oxadiazon | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Pendimethalin | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Permethrin | Insecticide | 0.030 | -- | -- | -- | -- | -- | -- | -- |
| Picloram | Herbicide | 0.05 | -- | -- | -- | -- | -- | 500 | 100 |
| Prometone | Herbicide | 0.05 | 1 | 1 | 0.9% | 0.0713 | -- | 100 | 20 |
| Prometryn | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Propiconazole | Fungicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Prothioconazole-desthio | Metabolite | 0.050 | -- | -- | -- | -- | -- | -- | -- |

| | | | | | | | | | |
|-----------------------|-------------|-------|----|----|-------|--------------|-------|-----|------|
| Saflufenacil | Herbicide | 0.05 | 2 | 6 | 5.7% | 0.0519-0.183 | 460 | -- | -- |
| Simazine | Herbicide | 0.05 | 2 | 4 | 3.8% | 0.0502-0.123 | -- | 4 | 0.4 |
| Sulfentrazone | Herbicide | 0.05 | 2 | 8 | 7.5% | 0.0633-0.45 | 1,000 | -- | -- |
| Sulfometuron methyl | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Tebupirimphos | Insecticide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Tembotrione | Herbicide | 0.10 | -- | -- | -- | -- | -- | -- | -- |
| Thiacloprid | Insecticide | 0.010 | -- | -- | -- | -- | -- | -- | -- |
| Thiamethoxam | Insecticide | 0.010 | 15 | 48 | 45.3% | 0.0316-3.54 | 120 | -- | -- |
| Thiencarbazone methyl | Herbicide | 0.05 | -- | -- | -- | -- | 800 | -- | -- |
| Triclopyr | Herbicide | 0.05 | -- | -- | -- | -- | -- | -- | -- |
| Trifluralin | Herbicide | 0.05 | -- | -- | -- | -- | -- | 7.5 | 0.75 |

Notes:

- 1 Total number of sites were 24.
 - 2 Total number of samples were 106.
 - 3 Wisconsin Department of Health Services(DHS) Drinking Water Health Advisory (June 2019, November 2020, revised February 2022).
 - 4 Combined sum of acetochlor metabolites ESA and OA.
 - 5 Combined sum of metabolites (di- and mono-acid) and parent material dacthal.
 - 6 Combined sum of metabolite DKN and parent material isoxaflutole.
 - 7 Combined sum of metolachlor metabolites ESA and OA.
 - Indicates that Health Advisory Level value in Wisconsin not established.
- DKN diketonitrile
 ESA ethane sulfonic acid
 OA oxanilic acid; can also be identified as OXA.
 µg/L. micrograms per liter or parts per billion
 TCR Total Chlorinated Residue for Atrazine. Reflects an additive quantity of atrazine (parent material) and its three metabolites (de-ethyl, de-isopropyl and di-amino atrazine).

| |
|--|
| |
| |
| |
| |

- Indicates no detects in excess of laboratory reporting limits.
- Indicates detects in excess of laboratory reporting limits.
- Indicates detects in excess of laboratory reporting limits and Wis. Admin. Code ch. NR 140 Preventive Action Limit.
- Indicates detects in excess of laboratory reporting limits and either Wis. Admin. Code ch. NR 140 Enforcement Standard or DHS Drinking Water Health Advisory.

Table B 4: Field-Edge Groundwater Monitoring Program - 2021 Land Pesticide/Nitrogen- and Irrigation-Use (as Provided by Growers)

| COUNTY | SITE (Grower) | YEAR | CROP | NUTRIENT MANAGEMENT PLAN | IRRIGATION APPLIED (in inches) | NITROGEN APPLIED (in lbs/acre) | PESTICIDE PRODUCT APPLIED | |
|-------------------|---------------|-------------------|-------------|--------------------------|--------------------------------|--------------------------------|--|-----|
| Adams | AD2 | 2016 | corn silage | --- | 6.45 | 374.8 | glyphosate atrazine dicamba | |
| | | 2017 ¹ | --- | --- | --- | --- | --- | |
| | | 2018 ¹ | --- | --- | --- | --- | --- | |
| | | 2019 ¹ | --- | --- | --- | --- | --- | |
| | | 2020 ¹ | --- | --- | --- | --- | --- | |
| | | 2021 ¹ | --- | --- | --- | --- | --- | |
| | AD3 | 2016 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2017 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2018 | snap beans | yes | 6.59 | 89.0 | metolachlor halosulfuron-methyl sethoxydim imazamox, bentazon thiamethoxam bifenthrin glyphosate | |
| | | 2019 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2020 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2021 ¹ | --- | --- | --- | --- | --- | --- |
| | AD4 | 2016 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2017 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2018 | soybeans | yes | 7.66 | 14.0 | metribuzin metolachlor clethodim bentazon thiamethoxam chlothianidin glyphosate | |
| | | 2019 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2020 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2021 ¹ | --- | --- | --- | --- | --- | --- |
| | AD5 | 2016 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2017 ¹ | --- | --- | --- | --- | --- | --- |
| | | 2018 ¹ | --- | --- | --- | --- | --- | --- |
| 2019 ¹ | | --- | --- | --- | --- | --- | --- | |
| 2020 ¹ | | --- | --- | --- | --- | --- | --- | |
| 2021 ¹ | | --- | --- | --- | --- | --- | --- | |
| Barron | BR3 | 2016 ¹ | --- | --- | --- | --- | --- | |
| | | 2017 ¹ | --- | --- | --- | --- | --- | |
| | | 2018 ¹ | --- | --- | --- | --- | --- | |
| | | 2019 | corn | no | 2.24 | 300 | glyphosate topramezone, dimethenamid acetochlor, flumetsulam, clopyralid | |
| | | 2020 ¹ | --- | --- | --- | --- | --- | |
| 2021 ¹ | --- | --- | --- | --- | --- | | | |
| Dane | DN1 | 2016 | seed corn | --- | 3 | 216.7 | simazine metolachlor mesotrione topramezone bifenthrin pyraclastrobin, metconazole 2,4-D glyphosate sodium chlorate | |
| | | 2017 | soybeans | --- | 2 | 6.0 | glyphosate clethodim lambda-cyhalothrin glufosinate | |
| | | 2018 ¹ | --- | --- | --- | --- | --- | |
| | | 2019 | soybeans | yes | 2 | 1.7 | glyphosate metribuzin dimethenamid glufosinate clethodim lambda-cyhalothrin | |
| | | 2020 | seed corn | yes | 4 | 201.95 | metolachlor glycine mesotrione simazine topramezone acetochlor simazine azoxystrobin, cyproconazole bifenthrin metaconazole, pyraclastrobin | |
| | | 2021 ¹ | --- | --- | --- | --- | --- | |
| | | 2016 | soybeans | --- | 3.43 | 100.0 | dimethenamid flumioxazin clethodim benzoic acid | |
| | | 2017 | horseradish | --- | 0.8 | 140.5 | peroxyacetic acid, hydrogen peroxide oxyfluorfen sulfentrazone glyphosate clethodim boscolid chlorothalonil glyphosate | |

| | | | | | | | | |
|------|-------|-------------------|-------------------|-----|------|--------|---|-----|
| Dunn | DU1 | 2018 | corn (grain) | no | 3.97 | 193.3 | dicamba dimethenamid, saflufenacil | |
| | | 2019 ¹ | --- | --- | --- | --- | --- | |
| | | 2020 | kidney beans | no | 2.5 | 91.98 | pendimethalin metolachlor imazamox sodium bentazon clethodim beta-cyfluthrin, imidacloprid saflufenacil | |
| | DU2 | 2021 | corn | no | 15.6 | 1076.9 | dicamba dimethenamide glyphosate saflufenacil glyphosate | |
| | | 2016 | corn | --- | 8 | 241.0 | dimethenamid, saflufenacil pendimethalin metolachlor bentazon | |
| | DU2 | 2017 | kidney beans | --- | 4 | 85.0 | fomesafen imazamox clethodim saflufenacil thiamethoxam, fludioxonil | |
| | | 2018 | corn | --- | 5 | 66.2 | dimethenamid, saflufenacil glyphosate atrazine pendimethalin glyphosate metolachlor imazamox | |
| | | 2019 | kidney beans | yes | 3.25 | 72.5 | bentazon fomesafen clethodim imidacloprid saflufenacil pendimethalin | |
| | | 2020 | kidney beans | no | 2.5 | 91.98 | metolachlor imazamox sodium bentazon clethodim beta-cyfluthrin, imidacloprid saflufenacil clothianidin | |
| | | 2021 | corn | no | 4.2 | 85 | glyphosate dicamba dimethenamide pyroxasulfone saflufenacil | |
| | Grant | GR1 | 2016 ¹ | --- | --- | na | --- | --- |
| | | | 2017 ¹ | --- | --- | na | --- | --- |
| | | | 2018 ¹ | --- | --- | na | --- | --- |
| | | | 2019 ¹ | --- | --- | na | --- | --- |
| | | | 2020 ¹ | --- | --- | na | --- | --- |
| IW1 | IW1 | 2021 ¹ | --- | --- | na | --- | --- | |
| | | 2016 | potatoes | --- | 18.4 | 374.4 | metam sodium azoxystrobin, difenoconazole metolachlor imidacloprid azoxystrobin metribuzin novaluron spinosad beta-cyfluthrin rimsulfuron chlorothalonil pyraclostrobin boscolid abamectin pyrimethanil mancozeb diquat bromide glyphosate | |
| | | 2017 | seed corn | --- | 8.9 | 198.5 | bifenthrin glufosinate MCPA, bromoxynil pendimethalin pyraclostrobin, metconazole propiconazole, azoxystrobin thiamethoxam | |
| | | 2018 | snap beans | no | 5.7 | 77.0 | halosulfuron-methyl s-metolachlor imazamox, bentazon sethoxydim | |
| | | 2019 ¹ | --- | --- | --- | --- | --- | |
| | | 2020 | potatoes | no | 21 | 225.93 | bifenthrin, pyraclostrobin metribuzin metolachlor indoxacarb acetamiprid chlorothalonil spinosad lambda-cyhalothrin mefentrifluconazole | |

| | | | | | | | |
|-------------------|-----|-------------------|-------------|-----|------|--------|--|
| Iowa | IW2 | 2021 | seed corn | no | 9.4 | 199 | abamectin zoxamide pyrimethanil mancozeb fentin hydroxide diquat dibromide abamectin azoxystrobin bifenthrin bromoxynil fludioxonil tembotrione glyphosate mefanoxam pendimethalin propiconazole pydiflumetofen thiabendazole thiamethoxam |
| | | 2016 | seed corn | --- | 12.8 | 195.5 | glyphosate bifenthrin metolachlor pendimethalin tembotrione bromoxynil azoxystrobin glyphosate |
| | | 2017 | snap beans | --- | 6.6 | 72.2 | EPTC thiamethoxam bifenthrin imazamox, bentazon |
| | | 2018 | seed corn | no | 12.1 | 256.0 | bifenthrin bicyclopyrone, metolachlor, mesotrione pendimethalin thiamethoxam azoxystrobin |
| | | 2019 ¹ | --- | --- | --- | --- | --- |
| | | 2020 | seed corn | no | 10.6 | 223.2 | bifenthrin glufosinate metolachlor nicosulfuron pyroxasulfone pendimethalin azoxystrobin, propiconazole, pydiflumetofen |
| | | 2021 | snap beans | no | 5.2 | 65 | bifenthrin captan glyphosate imazamox, bentazon halosulfuron-methyl matalaxyl sethoxydim metolachlor thiophanate-methyl thiram thiamethoxam |
| Jackson | JK3 | 2016 ¹ | --- | --- | na | --- | --- |
| | | 2017 ¹ | --- | --- | na | --- | --- |
| | | 2018 ¹ | --- | --- | na | --- | --- |
| | | 2019 ¹ | --- | --- | na | --- | --- |
| | | 2020 ¹ | --- | --- | na | --- | --- |
| | | 2021 ¹ | --- | --- | na | --- | --- |
| Juneau | JN1 | 2016 | sweet corn | --- | 8 | 211.0 | atrazine metolachlor |
| | | 2017 | snap beans | --- | 2.9 | 122.0 | metolachlor halosulfuron-methyl |
| | | 2018 | sweet corn | no | 8 | 228.6 | atrazine metolachlor |
| | | 2019 | potatoes | no | 12.5 | 65.05 | azoxystrobin chlorothalonil esfenvalerate spinosad thiamethoxam diquat dibromide boscalid metribuzin |
| | | 2020 | sweet corn | no | 9.5 | 212.37 | atrazine metolachlor |
| | | 2021 | snap beans | no | 5 | 152.6 | halosulfuron-methyl metolachlor |
| | JN3 | 2016 ¹ | --- | --- | na | --- | --- |
| | | 2017 ¹ | --- | --- | na | --- | --- |
| | | 2018 ¹ | --- | --- | na | --- | --- |
| | | 2019 ¹ | --- | --- | na | --- | --- |
| | | 2020 ¹ | --- | --- | na | --- | --- |
| 2021 ¹ | --- | --- | na | --- | --- | | |
| | | 2016 | corn silage | --- | --- | 179.5 | glyphosate lorsban acetochlor dicamba |

| | | | | | | | | |
|-------------|-----|-------------------|------------|-----|-------|--------|--|-----|
| La Crosse | LC2 | 2017 | soybeans | --- | --- | 0.0 | glyphosate 2,4-D imazethapyr | |
| | | 2018 | corn | yes | 2.5 | 705.7 | glyphosate atrazine, acetochlor mesotrione | |
| | | 2019 | beans | --- | --- | 0.0 | glyphosate methansulfonamide metribuzin metolachlor glyphosate, imazethapyr | |
| | | 2020 ¹ | --- | --- | --- | --- | --- | |
| | | 2021 ¹ | --- | --- | --- | --- | --- | |
| | | 2016 ¹ | --- | --- | --- | --- | --- | --- |
| Langlade | LN1 | 2017 ¹ | --- | --- | --- | --- | --- | |
| | | 2018 ¹ | --- | --- | --- | --- | --- | |
| | | 2019 ¹ | --- | --- | --- | --- | --- | |
| | | 2020 ¹ | --- | --- | --- | --- | --- | |
| | | 2021 ¹ | --- | --- | --- | --- | --- | |
| | | 2016 ¹ | --- | --- | --- | --- | --- | --- |
| Portage | PR1 | 2017 ¹ | --- | --- | --- | --- | --- | |
| | | 2018 | sweet corn | yes | 4.6 | 164.0 | metolachlor atrazine chlorothalonil azoxystrobin spinetram | |
| | | 2019 | potatoes | yes | 6.7 | 159 | abamectin, cyantraniliprole imidacloprid novaluron diqist | |
| | | 2020 ¹ | field corn | --- | 7.2 | 167.17 | glyphosate | |
| | | 2021 ¹ | --- | --- | --- | --- | --- | |
| | | 2016 ¹ | --- | --- | --- | --- | --- | --- |
| St. Croix | SC1 | 2016 | soybeans | --- | na | --- | glyphosate glyphosate tembotrione acetochlor glyphosate | |
| | | 2017 | corn | --- | na | 250.0 | --- | |
| | | 2018 | soybeans | no | na | 0.0 | --- | |
| | | 2019 ¹ | --- | --- | na | --- | --- | |
| | | 2020 ¹ | --- | --- | na | --- | --- | |
| | | 2021 ¹ | --- | --- | na | --- | --- | |
| Sauk | SK6 | 2016 ¹ | --- | --- | na | --- | --- | |
| | | 2017 ¹ | --- | --- | na | --- | --- | |
| | | 2018 ¹ | --- | --- | na | --- | --- | |
| | | 2019 ¹ | --- | --- | --- | --- | --- | |
| | | 2020 ¹ | --- | --- | --- | --- | --- | |
| | | 2021 ¹ | --- | --- | --- | --- | --- | |
| Trempealeau | TR1 | 2016 ¹ | --- | --- | --- | --- | --- | |
| | | 2017 ¹ | --- | --- | --- | --- | --- | |
| | | 2018 ¹ | --- | --- | --- | --- | --- | |
| | | 2019 ¹ | --- | --- | --- | --- | --- | |
| | | 2020 ¹ | --- | --- | --- | --- | --- | |
| | | 2021 ¹ | --- | --- | --- | --- | --- | |
| Waupaca | WP2 | 2016 | corn | --- | na | 132.0 | acetochlor clopyralid flumetsulam | |
| | | 2017 | soybeans | --- | na | 0.0 | glyphosate | |
| | | 2018 | soybeans | yes | na | 0.0 | glyphosate | |
| | | 2019 | corn | yes | na | 122.0 | acetochlor, clopyralid, flumetsulam glyphosate | |
| | | 2020 | corn | yes | na | 97.9 | acetochlor, clopyralid, flumetsulam | |
| | | 2021 | soybeans | yes | na | 0 | glyphosate | |
| | WS4 | 2016 | carrots | --- | 9.08 | 176.0 | glyphosate pendimethalin chlorothalonil esfenvalerate clethodim azoxystrobin glyphosate thiamethoxam, fludioxonil mancozeb azoxystrobin pentachloronitrobenzene metolachlor metribuzin rimsulfuron chlorothalonil novaluron metaxyl spinosad boscolid cyantraniliprole, abamectin pyraclostrobin oxathiapiprolin fentin hydroxide diqat bromide | |
| | | 2017 | potatoes | --- | 13.62 | 115.1 | metolachlor simazine glyphosate ammonium sulfamate metolachlor halosulfuron-methyl | |
| | | 2018 | corn | no | 9.1 | 70.6 | --- | |
| | | 2019 | beans | no | 2.42 | 24.96 | --- | |
| | | 2016 | --- | --- | --- | --- | --- | --- |
| | | 2017 | --- | --- | --- | --- | --- | --- |
| | | 2018 | --- | --- | --- | --- | --- | --- |
| | | 2019 | --- | --- | --- | --- | --- | --- |
| | | 2020 | --- | --- | --- | --- | --- | --- |
| | | 2021 | --- | --- | --- | --- | --- | --- |
| | | 2016 | --- | --- | --- | --- | --- | --- |

| | | | | | | | |
|----------|-----|------|----------|-----|-------|-----------------------------|---------------------|
| Waushara | | 2020 | carrots | no | 12.12 | 241.3 | pendimethalin |
| | | | | | | | clethodim |
| | | | | | | | prometryn |
| | | | | | | | carfentrazone-ethyl |
| | | | | | | | esfenvalerate |
| | | | | | | | chlorothalonil |
| | | | | | | | azoxystrobin |
| | | | | | | | boscalid |
| | | | | | | | abamectin |
| | | | | | | | cyantraniliprole |
| | | | | | | | esfenvalerate |
| | | | | | | | metolachlor |
| | | | | | | | novaluron |
| | | | | | | | pendimethalin |
| | | | | | | | phosmet |
| | | | | | | | spinetoram |
| | | | | | | | glyphosate |
| | | | | | | | simazine |
| | | | | | | | metolachlor |
| | | | | | | | glyphosate |
| | | | | | | | metolachlor |
| | | | | | | halosulfuron-methyl | |
| | | | | | | clethodim | |
| | | | | | | carfentrazone-ethyl | |
| | | | | | | cypermethrin | |
| | | | | | | azoxystrobin | |
| | | | | | | pendimethalin | |
| | | | | | | metribuzin | |
| | | | | | | novaluron | |
| | | | | | | phosmet | |
| | | | | | | chlorothalonil | |
| | | | | | | boscalid | |
| | | | | | | cyantraniliprole, abamectin | |
| | | | | | | metaxyl | |
| | | | | | | fentin hydroxide | |
| | | | | | | diquat dibromide | |
| | | | | | | glyphosate | |
| | | | | | | metolachlor | |
| | | | | | | simazine | |
| | | | | | | tembotrione | |
| | | | | | | metolachlor | |
| | | | | | | simazine | |
| | | | | | | topramezone | |
| | W56 | 2016 | corn | --- | 8.35 | 70.4 | |
| | | 2017 | beans | --- | 6 | 105.6 | |
| | | 2018 | carrots | no | 12.76 | 254.1 | |
| | | 2019 | potatoes | no | 10.9 | 200.16 | |
| | | 2020 | corn | no | 7.93 | 70.78 | |
| | | 2021 | corn | no | 14.6 | 133 | |
| | W57 | 2016 | | | | | |
| | | 2017 | | | | | |
| | | 2018 | | | | | |
| | | 2019 | | | | | |
| | | 2020 | | | | | |
| | | 2021 | | | | | |

Notes:

- 1 Grower did not provide information in Annual Reporting Form.
- Site is located within an atrazine Prohibition Area.
- Information not provided by Grower.
- na Fields are not equipped to irrigate.
- Site is a research location with multiple crops and herbicide types and application rates.

Table B 5: Field-Edge Groundwater Monitoring Program - 2021 Imidacloprid Concentrations in Groundwater Samples

| County | Site (Grower) | Well Name | WUWN | Sample Date | Imidacloprid | |
|-----------|---------------|-----------|------------|-------------|--------------|-----------|
| Adams | AD2 | AD2-1 | BH954 | 5/13/2021 | 0.0196 | |
| | | | | 11/24/2021 | 0.0167 | |
| | | AD2-4 | VR844 | 5/13/2021 | 2.77 | |
| | | | | 11/24/2021 | 2.27 | |
| | | AD2-5 | VR845 | 5/13/2021 | 0.356 | |
| | | | | 11/24/2021 | 0.31 | |
| | AD3 | AD2-6 | PT421 | | 11/24/2021 | 0 |
| | | AD3-1 | BH999 | | 5/13/2021 | 0.131 |
| | | AD3-3 | BI001 | | 5/13/2021 | 0.0812 |
| | AD4 | AD4-2 | BH997 | | 5/13/2021 | 0.143 |
| | | AD5 | AD5-1 | CL461 | | 5/13/2021 |
| | | | | | 11/24/2021 | 0 |
| | AD5-4 | | VR846 | | 5/13/2021 | 0.119 |
| | | | | | 11/24/2021 | 0.105 |
| AD5-5 | VR847 | | | 5/13/2021 | 0.299 | |
| | | | | 11/24/2021 | 0.326 | |
| | | AD5-6 | PT422 | | 11/24/2021 | 0 |
| Barron | BR3 | BR3-1 | BR279 | | 4/29/2021 | 0 |
| | | | | | 10/27/2021 | 0 |
| | | BR3-3 | BR281 | | 4/29/2021 | 0 |
| | | | | | 10/27/2021 | 0 |
| Dane | DN1 | DN1-1 | PT428 | | 10/21/2021 | 0 |
| | | DN1-2 | BR251 | | 5/6/2021 | 0 |
| | | DN1-3 | BR252 | | 10/21/2021 | 0.023 |
| Dunn | DU1 | DU1-1 | AO384 | | 5/25/2021 | 0 |
| | | | | | 10/27/2021 | 0 |
| | | DU1-3 | AO386 | | 5/25/2021 | 0 |
| | DU2 | | | | 10/27/2021 | 0 |
| | | DU2-1 | AO387 | | 5/25/2021 | 0 |
| | | DU2-3 | AO389 | | 10/27/2021 | 0 |
| Grant | GR1 | GR1-1 | BR255 | | 5/11/2021 | 0 |
| | | | | | 10/21/2021 | 0 |
| | | GR1-3 | BR257 | | 5/11/2021 | 0 |
| | | | | | 10/21/2021 | 0 |
| Iowa | IW1 | IW1-4 | BR259 | | 5/6/2021 | 0.142 |
| | | | | | 11/16/2021 | 0.138 |
| | | IW1-6 | BR261 | | 11/16/2021 | 0.0567 |
| | | IW1-7 | BH967 | | 5/6/2021 | 0.0172 |
| | | | | | 11/16/2021 | 0.0198 |
| | | IW1-8 | PT425 | | 11/16/2021 | 0.163 |
| | IW2 | IW2-1 | BR036 | | 5/6/2021 | 0 |
| | | IW2-2 | BR037 | | 11/16/2021 | 0.204 |
| | | IW2-3 | BR038 | | 5/6/2021 | 0.188 |
| | | IW2-4 | PT426 | | 11/16/2021 | 0.0274 |
| | | IW2-5 | PT427 | | 11/16/2021 | 0.0183 |
| Jackson | JK3 | JK3-1 | JH982 | | 5/4/2021 | 0 |
| | | | | | 10/27/2021 | 0 |
| | | JK3-2 | JH981 | | 5/4/2021 | 0 |
| | | | | | 10/27/2021 | 0 |
| Juneau | JN1 | JN1-1 | BR046 | | 4/7/2021 | 0 |
| | | | | | 12/2/2021 | 0.0177 |
| | | JN1-3 | BR048 | | 4/7/2021 | 0.0829 |
| | JN3 | | | | 12/2/2021 | 0.0833 |
| | | JN3-1 | JH937 | | 4/14/2021 | 0 |
| | | JN3-2 | JH936 | | 12/2/2021 | 0 |
| La Crosse | LC2 | LC2-1 | VZ391 | | 5/20/2021 | 0 |
| | | | | | 10/28/2021 | 0 |
| | | LC2-2 | VZ392 | | 5/20/2021 | 0 |
| | | | | | 10/28/2021 | 0 |
| Langlade | LN1 | LN1-1 | BH964 | | 5/18/2021 | 0.0201 |
| | | | | | 10/19/2021 | 0.017 |
| | | LN1-3 | BH966 | | 5/18/2021 | 0.0106 |
| | | | | | 10/19/2021 | 0 |
| Portage | PR1 | PR1-1 | BR207 | | 4/13/2021 | 0.0706 |
| | | | | | 10/19/2021 | 0 |
| | | PR1-4 | VR848 | | 4/13/2021 | 0.0428 |
| | | | | | 10/19/2021 | 0.034 |
| | | PR1-5 | VR849 | | 4/13/2021 | 0.0395 |
| | | | 10/19/2021 | 0.0363 | | |
| St. Croix | SC1 | SC1-1 | JH938 | | 4/29/2021 | 0 |
| | | | | | 10/28/2021 | 0 |
| | | SC1-2 | JH939 | | 4/29/2021 | 0 |
| | | | | | 10/28/2021 | 0 |

| | | | | | |
|-------------|-----|-------|-------|------------|--------|
| Sauk | SK6 | SK6-1 | BB246 | 5/11/2021 | 0.146 |
| | | SK6-2 | BB247 | 10/21/2021 | 0.272 |
| | | SK6-3 | BB248 | 5/11/2021 | 0.27 |
| | | SK6-4 | PT424 | 10/21/2021 | 0 |
| Trempealeau | TR1 | TR1-1 | PX201 | 5/20/2021 | 0 |
| | | | | 10/28/2021 | 0 |
| | | TR1-2 | PX202 | 5/20/2021 | 0 |
| | | | | 10/28/2021 | 0 |
| Waupaca | WP2 | WP2-1 | JH985 | 5/18/2021 | 0 |
| | | | | 10/19/2021 | 0 |
| | | WP2-2 | JH984 | 5/18/2021 | 0 |
| | | | | 10/19/2021 | 0 |
| Waushara | WS4 | WS4-1 | BB258 | 4/22/2021 | 0.192 |
| | | | | 11/3/2021 | 0.196 |
| | | WS4-4 | BB261 | 4/22/2021 | 0.0298 |
| | | | | 11/3/2021 | 0.028 |
| | WS6 | WS6-1 | JH989 | 4/22/2021 | 0.185 |
| | | | | 11/3/2021 | 0.0636 |
| | | WS6-2 | JH990 | 4/22/2021 | 0 |
| | | | | 11/3/2021 | 0 |
| | WS7 | WS7-1 | VR841 | 4/22/2021 | 0.0568 |
| | | | | 11/3/2021 | 0.0505 |
| | | WS7-2 | VR842 | 4/22/2021 | 0.261 |
| | | | | 11/3/2021 | 0.246 |
| | | WS7-3 | VR843 | 4/22/2021 | 0.84 |
| | | | | 11/3/2021 | 0.189 |
| | | WS7-4 | PT423 | 11/3/2021 | 0.0829 |

Notes:

WUWN
µg/L
0

Wisconsin Unique Well Number
Micrograms per liter or parts per billion
Concentration does not exceed laboratory reporting limit of 0.05 µg/L.
Exceeds Wisconsin Department of Health Services Drinking Water Health Advisory of 0.2 µg/L.
(June 2019, November 2020, revised February 2022)

Table B 6: Field-Edge Groundwater Monitoring Program - 2021 Alachlor ESA Concentrations in Groundwater Samples

| County | Site (Grower) | Well Name | WUWN | Sample Date | Alachlor ESA | | |
|-----------|---------------|------------|------------|-------------|--------------|------------|--------|
| Adams | AD2 | AD2-1 | BH954 | 5/13/2021 | 0.338 | | |
| | | | | 11/24/2021 | 0.14 | | |
| | | AD2-4 | VR844 | 5/13/2021 | 0.415 | | |
| | | | | 11/24/2021 | 0.411 | | |
| | | AD2-5 | VR845 | 5/13/2021 | 0.658 | | |
| | | | | 11/24/2021 | 0.528 | | |
| | AD2-6 | PT421 | 11/24/2021 | 3.31 | | | |
| | | | | | | | |
| | AD3 | AD3-1 | BH999 | 5/13/2021 | 1.1 | | |
| | | | | 5/13/2021 | 0.293 | | |
| | AD4 | AD4-2 | BH997 | 5/13/2021 | 0.182 | | |
| | | | | 5/13/2021 | 0 | | |
| AD5 | AD5-1 | CL461 | 11/24/2021 | 0 | | | |
| | | | 5/13/2021 | 1.36 | | | |
| | AD5-4 | VR846 | 11/24/2021 | 1.15 | | | |
| | | | 5/13/2021 | 11.3 | | | |
| | AD5-5 | VR847 | 11/24/2021 | 9.36 | | | |
| | | | | | | | |
| AD5-6 | PT422 | 11/24/2021 | 2.18 | | | | |
| | | | | | | | |
| Barron | BR3 | BR3-1 | BR279 | 4/29/2021 | 0 | | |
| | | | | 10/27/2021 | 0 | | |
| | | BR3-3 | BR281 | 4/29/2021 | 0 | | |
| | | | | 10/27/2021 | 0 | | |
| Dane | DN1 | DN1-1 | PT428 | 10/21/2021 | 0 | | |
| | | | | DN1-2 | BR251 | 5/6/2021 | 0.0611 |
| | | | | | | 10/21/2021 | 0 |
| Dunn | DU1 | DU1-1 | AO384 | 5/25/2021 | 0.188 | | |
| | | | | 10/27/2021 | 0.162 | | |
| | | DU1-3 | AO386 | 5/25/2021 | 0.166 | | |
| | 10/27/2021 | | | 0.137 | | | |
| | DU2 | DU2-1 | AO387 | 5/25/2021 | 0.094 | | |
| | | | | 10/27/2021 | 0.136 | | |
| DU2-3 | | AO389 | 5/25/2021 | 0.0816 | | | |
| | 10/27/2021 | | 0.0882 | | | | |
| Grant | GR1 | GR1-1 | BR255 | 5/11/2021 | 0.0799 | | |
| | | | | 10/21/2021 | 0 | | |
| | | GR1-3 | BR257 | 5/11/2021 | 0 | | |
| | | | | 10/21/2021 | 0.0728 | | |
| Iowa | IW1 | IW1-4 | BR259 | 5/6/2021 | 0.7 | | |
| | | | | 11/16/2021 | 1.12 | | |
| | | IW1-6 | BR261 | 11/16/2021 | 0.714 | | |
| | | | | 5/6/2021 | 1.22 | | |
| | | IW1-7 | BH967 | 11/16/2021 | 1.44 | | |
| | 11/16/2021 | | | 1.75 | | | |
| | IW2 | IW2-1 | BR036 | 5/6/2021 | 0.396 | | |
| | | | | 11/16/2021 | 0.271 | | |
| | | IW2-2 | BR037 | 5/6/2021 | 0.456 | | |
| | | | | 11/16/2021 | 0.359 | | |
| IW2-4 | | PT426 | 11/16/2021 | 0.438 | | | |
| | 11/16/2021 | | 0.351 | | | | |
| Iowa | IW2 | IW2-3 | BR038 | 5/6/2021 | 0.456 | | |
| | | | | 11/16/2021 | 0.359 | | |
| | | IW2-4 | PT426 | 11/16/2021 | 0.438 | | |
| | | | | 11/16/2021 | 0.351 | | |
| Jackson | JK3 | JK3-1 | JH982 | 5/4/2021 | 0 | | |
| | | | | 10/27/2021 | 0 | | |
| | | JK3-2 | JH981 | 5/4/2021 | 0 | | |
| | | | | 10/27/2021 | 0 | | |
| Juneau | JN1 | JN1-1 | BR046 | 4/7/2021 | 0 | | |
| | | | | 12/2/2021 | 0 | | |
| | | JN1-3 | BR048 | 4/7/2021 | 0.58 | | |
| | 12/2/2021 | | | 1.13 | | | |
| | JN3 | JN3-1 | JH937 | 4/14/2021 | 0 | | |
| | | | | 12/2/2021 | 15.5 | | |
| JN3-2 | | JH936 | 4/14/2021 | 0 | | | |
| | 12/2/2021 | | 0.11 | | | | |
| La Crosse | LC2 | LC2-1 | VZ391 | 5/20/2021 | 0 | | |
| | | | | 10/28/2021 | 0 | | |
| | | LC2-2 | VZ392 | 5/20/2021 | 0 | | |
| | | | | 10/28/2021 | 0 | | |
| Langlade | LN1 | LN1-1 | BH964 | 5/18/2021 | 0 | | |
| | | | | 10/19/2021 | 0 | | |
| | | LN1-3 | BH966 | 5/18/2021 | 0 | | |
| | | | | 10/19/2021 | 0 | | |
| Portage | PR1 | PR1-1 | BR207 | 4/13/2021 | 0 | | |
| | | | | 10/19/2021 | 0 | | |
| | | PR1-4 | VR848 | 4/13/2021 | 0.726 | | |
| | | | | 10/19/2021 | 0.656 | | |
| | | PR1-5 | VR849 | 4/13/2021 | 0.724 | | |
| | | | | 10/19/2021 | 0.722 | | |
| St. Croix | SC1 | SC1-1 | JH938 | 4/29/2021 | 0.245 | | |
| | | | | 10/28/2021 | 0.268 | | |
| | | SC1-2 | JH939 | 4/29/2021 | 0 | | |
| | | | | 10/28/2021 | 0.162 | | |

| | | | | | |
|-------------|-----------|-------|-----------|------------|-------|
| Sauk | SK6 | SK6-1 | BB246 | 5/11/2021 | 0.907 |
| | | SK6-2 | BB247 | 10/21/2021 | 0.727 |
| | | SK6-3 | BB248 | 5/11/2021 | 0.359 |
| | | SK6-4 | PT424 | 10/21/2021 | 0.611 |
| Trempealeau | TR1 | TR1-1 | PX201 | 5/20/2021 | 0 |
| | | | | 10/28/2021 | 0 |
| | | TR1-2 | PX202 | 5/20/2021 | 0 |
| | | | | 10/28/2021 | 0 |
| Waupaca | WP2 | WP2-1 | JH985 | 5/18/2021 | 0.132 |
| | | | | 10/19/2021 | 0.113 |
| | | WP2-2 | JH984 | 5/18/2021 | 0 |
| | | | | 10/19/2021 | 0 |
| Waushara | WS4 | WS4-1 | BB258 | 4/22/2021 | 0.22 |
| | | | | 11/3/2021 | 0.787 |
| | | WS4-4 | BB261 | 4/22/2021 | 0.386 |
| | | | | 11/3/2021 | 0.146 |
| | WS6 | WS6-1 | JH989 | 4/22/2021 | 0.16 |
| | | | | 11/3/2021 | 0.267 |
| | | WS6-2 | JH990 | 4/22/2021 | 0 |
| | 11/3/2021 | | | 0 | |
| | WS7 | WS7-1 | VR841 | 4/22/2021 | 0.189 |
| | | | | 11/3/2021 | 0.317 |
| | | WS7-2 | VR842 | 4/22/2021 | 0.944 |
| | | | | 11/3/2021 | 0.774 |
| | | WS7-3 | VR843 | 4/22/2021 | 3.11 |
| | | | | 11/3/2021 | 2.92 |
| WS7-4 | | PT423 | 11/3/2021 | 5.67 | |

Notes:

WUWN
µg/L
0

Wisconsin Unique Well Number
Micrograms per liter or parts per billion
Concentration does not exceed laboratory reporting limit of 0.05 µg/L.
Detected concentration exceeds the Wisconsin Administrative Code ch. NR 140 Preventive Action Limit of 4.0 µg/L.

Table B 7: Field-Edge Groundwater Monitoring Program - 2021 Atrazine and Metabolite Concentrations in Groundwater Samples

| County | Site (Grower) | Well Name | WUWN | Sample Date | Atrazine | De-ethyl Atrazine | De-isopropyl Atrazine | Di-amino Atrazine | Atrazine TCR |
|-------------|---------------|------------|------------|-------------|----------|-------------------|-----------------------|-------------------|--------------|
| Adams | AD2 | AD2-1 | BH954 | 5/13/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 11/24/2021 | 0 | 0 | 0 | 0 | 0 |
| | | AD2-4 | VR844 | 5/13/2021 | 0.17 | 0.197 | 0 | 0 | 0.367 |
| | | | | 11/24/2021 | 0.181 | 0.335 | 0 | 0 | 0.516 |
| | | AD2-5 | VR845 | 5/13/2021 | 0.0941 | 0.217 | 0 | 0 | 0.3111 |
| | | | | 11/24/2021 | 0.179 | 0.25 | 0 | 0 | 0.429 |
| | AD2-6 | PT421 | 11/24/2021 | 0.328 | 0.728 | 0 | 0.241 | 1.297 | |
| | AD3 | AD3-1 | BH999 | 5/13/2021 | 0 | 0.581 | 0 | 0.229 | 0.81 |
| | | AD3-3 | BI001 | 5/13/2021 | 0 | 0 | 0 | 0 | 0 |
| | AD4 | AD4-2 | BH997 | 5/13/2021 | 0 | 0.0634 | 0 | 0 | 0.0634 |
| | | | | 11/24/2021 | 0 | 0 | 0 | 0 | 0 |
| | | AD5-1 | CL461 | 5/13/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 11/24/2021 | 0 | 0.0529 | 0.0935 | 0 | 0.1464 |
| | | AD5-4 | VR846 | 5/13/2021 | 0.0696 | 0.0595 | 0.0573 | 0 | 0.1864 |
| | | | | 11/24/2021 | 0.0603 | 0 | 0 | 0 | 0.0603 |
| | AD5-5 | VR847 | 5/13/2021 | 0.136 | 0.744 | 0 | 0.205 | 1.085 | |
| 11/24/2021 | | | 0.142 | 0.722 | 0 | 0 | 0.854 | | |
| AD5-6 | PT422 | 11/24/2021 | 0 | 0.606 | 0 | 0.201 | 0.807 | | |
| Barron | BR3 | BR3-1 | BR279 | 4/29/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/27/2021 | 0 | 0 | 0 | 0 | 0 |
| | | BR3-3 | BR281 | 4/29/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/27/2021 | 0 | 0 | 0 | 0 | 0 |
| Dane | DN1 | DN1-1 | PT428 | 10/21/2021 | 0 | 0 | 0 | 0 | 0 |
| | | DN1-2 | BR251 | 5/6/2021 | 0 | 0 | 0 | 0 | 0 |
| | | DN1-3 | BR252 | 10/21/2021 | 0 | 0 | 0 | 0 | 0 |
| Dunn | DU1 | DU1-1 | AO384 | 5/25/2021 | 0 | 0 | 0.141 | 0 | 0.141 |
| | | | | 10/27/2021 | 0.0797 | 0 | 0.113 | 0 | 0.1927 |
| | | | | 5/25/2021 | 0 | 0 | 0.227 | 0 | 0.227 |
| | DU1-3 | AO386 | 10/27/2021 | 0 | 0 | 0.195 | 0 | 0.195 | |
| | | | 5/25/2021 | 0 | 0 | 0 | 0 | 0 | |
| | | | 10/27/2021 | 0 | 0 | 0 | 0 | 0 | |
| DU2 | DU2-1 | AO387 | 5/25/2021 | 0 | 0 | 0 | 0 | 0 | |
| | | | 10/27/2021 | 0 | 0 | 0 | 0 | 0 | |
| | | | 5/25/2021 | 0 | 0 | 0 | 0 | 0 | |
| DU2-3 | AO389 | 10/27/2021 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | | 5/11/2021 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | | 10/21/2021 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Grant | GR1 | GR1-1 | BR255 | 5/11/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/21/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 5/11/2021 | 0 | 0 | 0 | 0 | 0 |
| Iowa | IW1 | IW1-4 | BR259 | 5/6/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 11/16/2021 | 0 | 0 | 0 | 0 | 0 |
| | | IW1-6 | BR261 | 11/16/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 5/6/2021 | 0 | 0.054 | 0.054 | 0 | 0.108 |
| | | IW1-7 | BH967 | 11/16/2021 | 0 | 0.0546 | 0.0556 | 0 | 0.1102 |
| | IW2 | IW2-1 | PT425 | 11/16/2021 | 0.089 | 0.101 | 0.17 | 0.261 | 0.621 |
| | | | | 5/6/2021 | 0 | 0 | 0 | 0 | 0 |
| | | IW2-2 | BR037 | 11/16/2021 | 0 | 0 | 0.0521 | 0 | 0.0521 |
| | | | | 5/6/2021 | 0 | 0 | 0.0507 | 0 | 0.0507 |
| | | IW2-3 | BR038 | 11/16/2021 | 0 | 0 | 0 | 0 | 0 |
| IW2-4 | PT426 | 11/16/2021 | 0 | 0 | 0.0702 | 0.247 | 0.3172 | | |
| | | 11/16/2021 | 0.172 | 0.102 | 0.065 | 0 | 0.339 | | |
| Jackson | JK3 | JK3-1 | JH982 | 5/4/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/27/2021 | 0 | 0 | 0 | 0 | 0 |
| | | JK3-2 | JH981 | 5/4/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/27/2021 | 0 | 0 | 0 | 0 | 0 |
| Juneau | JN1 | JN1-1 | BR046 | 4/7/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 12/2/2021 | 0 | 0 | 0 | 0 | 0 |
| | | JN1-3 | BR048 | 4/7/2021 | 0 | 0.0699 | 0 | 0 | 0.0699 |
| | JN3 | JN3-1 | JH937 | 4/14/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 12/2/2021 | 0 | 0 | 0 | 0 | 0 |
| | | JN3-2 | JH936 | 4/14/2021 | 0 | 0 | 0 | 0 | 0 |
| La Crosse | LC2 | LC2-1 | VZ391 | 5/20/2021 | 0 | 0.171 | 0 | 0 | 0.171 |
| | | | | 10/28/2021 | 0.0536 | 0.157 | 0 | 0 | 0.2106 |
| | | LC2-2 | VZ392 | 5/20/2021 | 0.0865 | 0.167 | 0 | 0 | 0.2535 |
| | | | | 10/28/2021 | 0.0672 | 0.165 | 0 | 0 | 0.2322 |
| Langlade | LN1 | LN1-1 | BH964 | 5/18/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/19/2021 | 0 | 0 | 0 | 0 | 0 |
| | | LN1-3 | BH966 | 5/18/2021 | 0 | 0 | 0 | 0 | 0 |
| Portage | PR1 | PR1-1 | BR207 | 4/13/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/19/2021 | 0 | 0 | 0 | 0 | 0 |
| | | PR1-4 | VR848 | 4/13/2021 | 0 | 0.0714 | 0 | 0 | 0.0714 |
| | | | | 10/19/2021 | 0 | 0.0729 | 0 | 0 | 0.0729 |
| | | PR1-5 | VR849 | 4/13/2021 | 0 | 0.0996 | 0 | 0 | 0.0996 |
| | | | | 10/19/2021 | 0 | 0.098 | 0 | 0 | 0.098 |
| St. Croix | SC1 | SC1-1 | JH938 | 4/29/2021 | 0 | 0.0636 | 0 | 0.247 | 0.3106 |
| | | | | 10/28/2021 | 0 | 0.0678 | 0.0511 | 0.23 | 0.3489 |
| | | SC1-2 | JH939 | 4/29/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/28/2021 | 0 | 0 | 0 | 0 | 0 |
| Sauk | SK6 | SK6-1 | BB246 | 5/11/2021 | 0 | 0 | 0 | 0 | 0 |
| | | SK6-2 | BB247 | 10/21/2021 | 0 | 0 | 0 | 0 | 0 |
| | | SK6-3 | BB248 | 5/11/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/21/2021 | 0 | 0 | 0 | 0 | 0 |
| | | SK6-4 | PT424 | 10/21/2021 | 0.116 | 0.272 | 0.503 | 0.382 | 1.273 |
| Trempealeau | TR1 | TR1-1 | PX201 | 5/20/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/28/2021 | 0 | 0 | 0 | 0 | 0 |
| | | TR1-2 | PX202 | 5/20/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 10/28/2021 | 0 | 0 | 0 | 0 | 0 |

| | | | | | | | | | |
|----------|-------|-------|-----------|------------|--------|--------|--------|-------|--------|
| Waupaca | WP2 | WP2-1 | JH985 | 5/18/2021 | 0 | 0.07 | 0.0729 | 0.253 | 0.3959 |
| | | | | 10/19/2021 | 0 | 0.0564 | 0.0773 | 0 | 0.1337 |
| | | WP2-2 | JH984 | 5/18/2021 | 0 | 0.0923 | 0 | 0 | 0.0923 |
| | | | | 10/19/2021 | 0 | 0.116 | 0.0716 | 0 | 0.1876 |
| Waushara | WS4 | WS4-1 | BB258 | 4/22/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 11/3/2021 | 0 | 0 | 0 | 0 | 0 |
| | | WS4-4 | BB261 | 4/22/2021 | 0 | 0 | 0.187 | 0.206 | 0.393 |
| | | | | 11/3/2021 | 0 | 0 | 0.145 | 0 | 0.145 |
| | WS6 | WS6-1 | JH989 | 4/22/2021 | 0 | 0 | 0.355 | 0.468 | 0.823 |
| | | | | 11/3/2021 | 0 | 0 | 0.605 | 0.512 | 1.117 |
| | WS7 | WS6-2 | JH990 | 4/22/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 11/3/2021 | 0 | 0 | 0 | 0 | 0 |
| | | WS7-1 | VR841 | 4/22/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 11/3/2021 | 0 | 0 | 0 | 0 | 0 |
| | | WS7-2 | VR842 | 4/22/2021 | 0 | 0 | 0 | 0 | 0 |
| | | | | 11/3/2021 | 0 | 0 | 0 | 0 | 0 |
| | | WS7-3 | VR843 | 4/22/2021 | 0.0641 | 0.426 | 0.269 | 0 | 0.7591 |
| | | | | 11/3/2021 | 0.0742 | 0.379 | 0.184 | 0 | 0.6372 |
| | WS7-4 | PT423 | 11/3/2021 | 0.104 | 0.495 | 0 | 0 | 0.599 | |

Concentrations identified as micrograms per liter or parts per billion.

TCR Total Chlorinated Residue for Atrazine. Reflects an additive quantity of atrazine and its three metabolites (de-ethyl, de-isopropyl and di-amino atrazine).

WUWN Wisconsin Unique Well Number

µg/L Micrograms per liter or parts per billion.

0 Concentration does not exceed laboratory reporting limit of 0.05 µg/L.

Site is located within an atrazine Prohibition Area.

Detected concentration exceeds the Wisconsin Administrative Code ch. NR 140 Preventive Action Limit of 0.3 µg/L.

Table B 8: Field-Edge Groundwater Monitoring Program - 2021 Nitrogen-Nitrate/Nitrite Concentrations in Groundwater Samples

| County | Site (Grower) | Well Name | WUWN | Sample Date | Nitrogen-Nitrate/Nitrite |
|------------|---------------|------------|------------|-------------|--------------------------|
| Adams | AD2 | AD2-1 | BH954 | 5/13/2021 | 29.6 |
| | | | | 11/24/2021 | 34 |
| | | AD2-4 | VR844 | 5/13/2021 | 33.1 |
| | | | | 11/24/2021 | 27.5 |
| | | AD2-5 | VR845 | 5/13/2021 | 28.8 |
| | | | | 11/24/2021 | 19.4 |
| | AD2-6 | PT421 | 11/24/2021 | 13 | |
| | | | | | |
| | AD3 | AD3-1 | BH999 | 5/13/2021 | 19.3 |
| | | | | AD3-3 | BI001 |
| | AD4 | AD4-2 | BH997 | 5/13/2021 | 31.4 |
| | | | | | |
| | AD5 | AD5-1 | CL461 | 5/13/2021 | 1.57 |
| | | | | 11/24/2021 | 1.43 |
| AD5-4 | | VR846 | 5/13/2021 | 20.2 | |
| | | | 11/24/2021 | 24.4 | |
| AD5-5 | | VR847 | 5/13/2021 | 30.2 | |
| | | | 11/24/2021 | 26.1 | |
| AD5-6 | PT422 | 11/24/2021 | 7.85 | | |
| Barron | BR3 | BR3-1 | BR279 | 4/29/2021 | 1.26 |
| | | | | 10/27/2021 | 2.67 |
| | | BR3-3 | BR281 | 4/29/2021 | 3.89 |
| | | | | 10/27/2021 | 9.83 |
| Dane | DN1 | DN1-1 | PT428 | 10/21/2021 | 10.8 |
| | | DN1-2 | BR251 | 5/6/2021 | 17 |
| | | DN1-3 | BR252 | 10/21/2021 | 14.5 |
| Dunn | DU1 | DU1-1 | AO384 | 5/25/2021 | 9.41 |
| | | | | 10/27/2021 | 12.4 |
| | | DU1-3 | AO386 | 5/25/2021 | 15.5 |
| | | | | 10/27/2021 | 12.6 |
| | DU2 | DU2-1 | AO387 | 5/25/2021 | 1.81 |
| | | | | 10/27/2021 | 6.85 |
| DU2-3 | AO389 | 5/25/2021 | 0 | | |
| | | 10/27/2021 | 0 | | |
| Grant | GR1 | GR1-1 | BR255 | 5/11/2021 | 14.1 |
| | | | | 10/21/2021 | 10.6 |
| | | | | 5/11/2021 | 18.1 |
| | | GR1-3 | BR257 | 10/21/2021 | 18.1 |
| | | | | | |
| Iowa | IW1 | IW1-4 | BR259 | 5/6/2021 | 16 |
| | | | | 11/16/2021 | 18.5 |
| | | IW1-6 | BR261 | 11/16/2021 | 13.3 |
| | | | | 5/6/2021 | 26.3 |
| | | IW1-7 | BH967 | 11/16/2021 | 25.8 |
| | 11/16/2021 | | | 26.5 | |
| | IW2 | IW2-1 | BR036 | 5/6/2021 | 0 |
| | | | | 11/16/2021 | 20.6 |
| | | | | 5/6/2021 | 28.2 |
| | | | | 11/16/2021 | 22.9 |
| 11/16/2021 | | | | 23.9 | |
| IW2-5 | PT427 | 11/16/2021 | 17 | | |
| | | | | | |
| Jackson | JK3 | JK3-1 | JH982 | 5/4/2021 | 2.4 |
| | | | | 10/27/2021 | 1.96 |
| | | JK3-2 | JH981 | 5/4/2021 | 2.72 |
| | | | | 10/27/2021 | 2.17 |
| Juneau | JN1 | JN1-1 | BR046 | 4/7/2021 | 5.72 |
| | | | | 12/2/2021 | 6.38 |
| | | JN1-3 | BR048 | 4/7/2021 | 31.1 |
| | | | | 12/2/2021 | 27 |
| | JN3 | JN3-1 | JH937 | 4/14/2021 | 0 |
| | | | | 12/2/2021 | 5.11 |
| JN3-2 | JH936 | 4/14/2021 | 1.31 | | |
| | | 12/2/2021 | 0 | | |
| La Crosse | LC2 | LC2-1 | VZ391 | 5/20/2021 | 19.1 |
| | | | | 10/28/2021 | 19.4 |
| | | LC2-2 | VZ392 | 5/20/2021 | 17.2 |
| | | | | 10/28/2021 | 17.6 |
| Langlade | LN1 | LN1-1 | BH964 | 5/18/2021 | 14.8 |
| | | | | 10/19/2021 | 15.5 |
| | | LN1-3 | BH966 | 5/18/2021 | 11.7 |
| | | | | 10/19/2021 | 10.6 |
| Portage | PR1 | PR1-1 | BR207 | 4/13/2021 | 4.9 |
| | | | | 10/19/2021 | 0.772 |
| | | PR1-4 | VR848 | 4/13/2021 | 21.6 |
| | | | | 10/19/2021 | 19.4 |
| | | PR1-5 | VR849 | 4/13/2021 | 22.6 |
| 10/19/2021 | 22.6 | | | | |

| | | | | | |
|-------------|-----|-----------|-----------|------------|------|
| St. Croix | SC1 | SC1-1 | JH938 | 4/29/2021 | 8.42 |
| | | | | 10/28/2021 | 10.2 |
| Sauk | SK6 | SC1-2 | JH939 | 4/29/2021 | 11.5 |
| | | SK6-1 | BB246 | 10/28/2021 | 19.4 |
| | | SK6-2 | BB247 | 5/11/2021 | 22.7 |
| | | SK6-3 | BB248 | 10/21/2021 | 41.2 |
| Trempealeau | TR1 | SK6-4 | PT424 | 5/11/2021 | 23.2 |
| | | | | 10/21/2021 | 17.3 |
| | | TR1-1 | PX201 | 10/21/2021 | 7.25 |
| | | TR1-2 | PX202 | 5/20/2021 | 25.5 |
| Waupaca | WP2 | | | 10/28/2021 | 23.6 |
| | | WP2-1 | JH985 | 5/20/2021 | 17.5 |
| | | WP2-2 | JH984 | 10/28/2021 | 17.7 |
| | | | | 5/18/2021 | 8.57 |
| Waushara | WS4 | WS4-1 | BB258 | 10/19/2021 | 32.6 |
| | | WS4-4 | BB261 | 5/18/2021 | 8.79 |
| | WS6 | WS6-1 | JH989 | 10/19/2021 | 13.3 |
| | | WS6-2 | JH990 | 4/22/2021 | 16.3 |
| | WS7 | WS7-1 | VR841 | 11/3/2021 | 18.6 |
| | | WS7-2 | VR842 | 4/22/2021 | 28.3 |
| | | WS7-3 | VR843 | 11/3/2021 | 25.8 |
| | | WS7-4 | PT423 | 4/22/2021 | 28.8 |
| | | | | 11/3/2021 | 34.2 |
| | | | | 4/22/2021 | 4.3 |
| | | | 11/3/2021 | 2.29 | |
| | | | 4/22/2021 | 14.3 | |
| | | | 11/3/2021 | 16.7 | |
| | | | 4/22/2021 | 31.9 | |
| | | 11/3/2021 | 23.3 | | |
| | | 4/22/2021 | 33.5 | | |
| | | 11/3/2021 | 31.4 | | |
| | | 11/3/2021 | 21.9 | | |

Notes:

| | |
|------|--|
| WUWN | Wisconsin Unique Well Number |
| mg/L | Milligrams per liter or parts per million |
| 0 | Concentration does not exceed laboratory reporting limit of 0.5 mg/L. |
| | Detected concentration exceeds the Wisconsin Administrative Code ch. NR 140 Preventive Action Limit of 2.0 mg/L. |
| | Detected concentration exceeds the Wisconsin Administrative Code ch. NR 140 Enforcement Standard of 10.0 mg/L. |